

Patient NAME :		Report STATUS : Final Report	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type : Whole blood EDTA	
Referred BY : Self		Report Date : Feb 26, 2026, 05:42 PM.	
Sample Collected : Feb 26, 2026, 11:46 AM			
Test Description	Value(s)	Unit(s)	Reference Range

Fever Package- Comprehensive

Complete Blood Count (CBC)

RBC Parameters			
Hemoglobin <i>colorimetric</i>	15.9	g/dL	13.0 - 17.0
RBC Count <i>Electrical impedance</i>	5.1	10 ⁶ /μl	4.5 - 5.5
PCV <i>Calculated</i>	45.1	%	40 - 50
MCV <i>Calculated</i>	87.7	fl	83 - 101
MCH <i>Calculated</i>	31	pg	27 - 32
MCHC <i>Calculated</i>	35.3 H*	g/dL	31.5 - 34.5
RDW (CV) * <i>Calculated</i>	13.3	%	11.6 - 14.0
RDW-SD * <i>Calculated</i>	42.8	fl	35.1 - 43.9
WBC Parameters			
TLC <i>Electrical impedance and microscopy</i>	3.4 L*	10 ³ /μl	4 - 10
Differential Leucocyte Count			
Neutrophils	41	%	40-80
Lymphocytes	48.6 H*	%	20-40
Monocytes	4.7	%	2-10
Eosinophils	4.3	%	1-6
Basophils	1.4	%	<2
Absolute Leukocyte Counts <i>calculated</i>			
Neutrophils.	1.39 L*	10 ³ /μl	2 - 7
Lymphocytes.	1.65	10 ³ /μl	1 - 3
Monocytes.	0.16 L*	10 ³ /μl	0.2 - 1.0
Eosinophils.	0.15	10 ³ /μl	0.02 - 0.5
Basophils.	0.05	10 ³ /μl	0.02 - 0.5
Platelet Parameters			
Platelet Count <i>Electrical impedance and microscopy</i>	148 L*	10 ³ /μl	150 - 410
The platelet count verified with peripheral smear and occasional small clumps with giant platelets seen. Manual platelet count is approximately around 1,50,000 to 1,60,000 per cu.mm			
Mean Platelet Volume (MPV) *	9.9	fL	9.3 - 12.1

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

K. Kokila

Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO :
Patient ID / UHID :		Sample Type : Whole blood EDTA
Referred BY : Self		Report Date : Feb 26, 2026, 05:42 PM.
Sample Collected : Feb 26, 2026, 11:46 AM		


Test Description	Value(s)	Unit(s)	Reference Range
<i>Calculated</i>			
PCT * <i>Calculated</i>	0.2	%	0.17 - 0.32
PDW * <i>Calculated</i>	17.3	fL	8.3 - 25.0
P-LCR * <i>Calculated</i>	30.9	%	18 - 50
P-LCC * <i>Calculated</i>	46	10 ⁹ /L	44 - 140
Mentzer Index * <i>Calculated</i>	17.2	%	> 13

Interpretation:

CBC provides information about red cells, white cells and platelets. Results are useful in the diagnosis of anemia, infections, leukemias, clotting disorders and many other medical conditions.

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

📞 898-898-0606

✉ care@redcliffelabs.com

🌐 www.redcliffelabs.com

All Lab results are subject to clinical interpretation by qualified medical professional and this report is not subject to use for any medico-legal purpose.

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO :
Patient ID / UHID :		Sample Type : Whole blood EDTA
Referred BY : Self		Report Date : Feb 26, 2026, 06:03 PM.
Sample Collected : Feb 26, 2026, 11:46 AM		

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Erythrocyte Sedimentation Rate (ESR)


ESR - Erythrocyte Sedimentation Rate <i>MODIFIED WESTERGREIN</i>	2	mm/hr	0 - 10
---	---	-------	--------

Interpretation:

ESR is also known as Erythrocyte Sedimentation Rate. An ESR test is used to assess inflammation in the body. Many conditions can cause an abnormal ESR, so an ESR test is typically used with other tests to diagnose and monitor different diseases. An elevated ESR may occur in inflammatory conditions including infection, rheumatoid arthritis, systemic vasculitis, anemia, multiple myeloma, etc. Low levels are typically seen in congestive heart failure, polycythemia, sickle cell anemia, hypo fibrinogenemia, etc.

Reference- Dacie and Lewis practical hematology

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO :
Patient ID / UHID :		Sample Type : Whole blood EDTA
Referred BY : Self		Report Date : Feb 26, 2026, 05:42 PM.
Sample Collected : Feb 26, 2026, 11:46 AM		

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------


Malarial Parasite (MP) Smear

MP(PBF FOR MP) MICROSCOPY	Not Seen		NOT SEEN
------------------------------	----------	--	----------

Interpretation:

1. Malaria is a serious parasitic diseases characterized by fever, chills, and anemia and is caused by a parasite that is transmitted human to human by the bite of infected female Anopheles mosquitoes.
2. Malarial Parasite test is performed on the blood sample to find out the level of Malaria Parasite in the blood.
3. It is conducted to conclude on Malaria and also during the treatment and after the treatment of Malaria.
4. Most people will have symptoms within 14 days of being bitten by an infected mosquito. But symptoms can show up as soon as seven days afterward or can take as long as a year to appear.
5. Clinical decision should not be based on the results of this test, but should be made by the physician after all clinical and laboratory findings have been evaluated.

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO
Patient ID / UHID :		Sample Type : Whole blood EDTA
Referred BY : Self		Report Date : Feb 26, 2026, 05:42 PM.
Sample Collected : Feb 26, 2026, 11:46 AM		

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------


Malaria Antigen, Rapid Card

Plasmodium Vivax	Negative		Negative
Plasmodium falciparum	Negative		Negative

Interpretation:

Immunochromatographic Assay done for Plasmodium falciparum using Histidine-Rich Protein-II (HRP-II) and Plasmodium species (Plasmodium falciparum, P. vivax, P. ovale and P. malariae) using lactate dehydrogenase (pLDH) in human whole blood.

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO
Patient ID / UHID :		Sample Type : Serum
Referred BY : Self		Report Date : Feb 26, 2026, 06:38 PM.
Sample Collected : Feb 26, 2026, 11:46 AM		

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Bilirubin (Total, Direct, Indirect)

Bilirubin Total <i>Photometric</i>	0.5	mg/dL	0.2 - 1.2
Bilirubin Direct * <i>Diazo Reaction</i>	0.3	mg/dL	0.0 - 0.5
Bilirubin Indirect * <i>Calculation (T Bil - D Bil)</i>	0.2	mg/dL	0.1 - 1.0

Interpretation:

Adults and children

Increased total bilirubin that is mainly unconjugated (indirect) bilirubin may be a result of:-

1. Hemolytic or pernicious anemia
2. Transfusion reaction
3. Cirrhosis
4. A relatively common inherited condition called Gilbert syndrome, due to low levels of the enzyme that produces conjugated bilirubin.

Newborns

An elevated bilirubin level in a newborn may be temporary and resolve itself within a few days to two weeks. However, if the bilirubin level is above a critical threshold or increases rapidly, an investigation of the cause is needed so appropriate treatment can be initiated. Increased bilirubin concentrations may result from the accelerated breakdown of red blood cells due to:

1. Blood type incompatibility between the mother and her newborn
2. Certain congenital infections
3. Lack of oxygen (hypoxia)
4. Diseases that can affect the liver

In most of these conditions, only unconjugated (indirect) bilirubin is increased.

SGOT / AST

SGOT/AST <i>IFCC without P5P</i>	37.4 H*	U/L	5 - 34
-------------------------------------	---------	-----	--------

Interpretation:

Serum AST is used for differential diagnosis of diseases of hepatobiliary system and pancreas. Increased values are seen in liver diseases like acute viral hepatitis, cirrhosis, biliary obstruction, primary or metastatic cancer, granuloma, hepatic ischaemia.

SGPT / ALT

SGPT/ALT <i>IFCC without P5P</i>	50.6	U/L	0 to 55
-------------------------------------	------	-----	---------

Interpretation:

Serum ALT is used for differential diagnosis of diseases of hepatobiliary system and pancreas. Increased in alcoholic hepatitis, cirrhosis, hepatocellular carcinoma, chronic hepatitis. Decreased in genito-urinary tract infection, malignancy, pyridoxal phosphate deficiency states (malnutrition, pregnancy, alcoholic liver disease).

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

K. Kokila

Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO
Patient ID / UHID :		Sample Type : Serum
Referred BY : Self		Report Date : Feb 26, 2026, 06:38 PM.
Sample Collected : Feb 26, 2026, 11:46 AM		

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Kidney Function Test (KFT)


Blood Urea <i>Urease</i>	25.4	mg/dL	19 - 44.1
Bun * <i>Urease</i>	11.87	mg/dL	8.9 - 20.6
Creatinine <i>Photometric</i>	1	mg/dL	0.72 - 1.25
eGFR (CKD-EPI)	97.56	ml/min/1.73 sq m	Normal Or High: >= 90 Mild Or Decrease: 60-89 Mild To Moderate Decrease: 45-59 Mild To Severe Decrease: 30-44 Severe Decrease: 15-29 Kidney Failure: < 15
Bun/Creatinine Ratio * <i>Calculated</i>	11.87 L*		12 - 20
Urea / Creatinine Ratio * <i>Calculated</i>	25.4 L*		25.68- 42.8
Uric Acid <i>Uricase</i>	5.4	mg/dL	3.5 - 7.2
Calcium Serum <i>Arsenazo III</i>	9	mg/dL	8.4 - 10.2
Phosphorus <i>Photometric</i>	4.3	mg/dL	2.3 - 4.7
Sodium <i>Potentiometric</i>	143.7	mmol/L	136 - 145
Potassium <i>Potentiometric</i>	3.7	mmol/L	3.5 - 5.1
Chloride <i>Potentiometric</i>	104	mmol/L	98 - 107

Interpretation:

Kidney function tests is a collective term for a variety of individual tests and procedures that can be done to evaluate how well the kidneys are functioning. Many conditions can affect the ability of the kidneys to carry out their vital functions. Some lead to a rapid (acute) decline in kidney function others lead to a gradual (chronic) decline in function. Both result in a buildup of toxic waste substances done on urine samples, as well as on blood samples. A number of symptoms may indicate a problem with your kidneys. These include : high blood pressure, blood in urine, frequent urges to urinate, difficulty beginning urination, painful urination, swelling in the hands and feet due to a buildup of fluids in the body. A single symptom may not mean something serious. However, when occurring simultaneously, these symptoms suggest that your kidneys are not working properly. Kidney function tests can help determine the reason. Ionized calcium this test if you have signs of kidney or parathyroid disease. The test may also be done to monitor progress and treatment of these diseases."eGFR test is applicable for patients aged 18 years or more."

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

☎ 898-898-0606

✉ care@redcliffelabs.com

🌐 www.redcliffelabs.com

All Lab results are subject to clinical interpretation by qualified medical professional and this report is not subject to use for any medico-legal purpose.

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO
Patient ID / UHID :		Sample Type : Serum
Referred BY : Self		Report Date : Feb 26, 2026, 06:38 PM.
Sample Collected : Feb 26, 2026, 11:46 AM		

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

C-Reactive Protein (CRP), Quantitative

CRP (Quantitative) <i>Immunoturbidimetry</i>	7.1 H*	mg/L	up to 5
---	--------	------	---------

Interpretation:


Increased CRP level:

1. A high or increasing amount of CRP in the blood suggests the presence of inflammation but will not identify its location or the cause.
2. Suspected bacterial infection—a high CRP level can provide indication that patient has an infection.
3. Chronic inflammatory disease—high levels of CRP suggest a flare-up if you have a chronic inflammatory disease or that treatment has not been effective.

If the CRP level is initially elevated and drops, it means that the inflammation or infection is subsiding and/or responding to treatment.

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

📞 898-898-0606

✉ care@redcliffelabs.com

🌐 www.redcliffelabs.com

All Lab results are subject to clinical interpretation by qualified medical professional and this report is not subject to use for any medico-legal purpose.

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO
Patient ID / UHID :		Sample Type : Serum
Referred BY : Self		Report Date : Feb 26, 2026, 08:27 PM.
Sample Collected : Feb 26, 2026, 11:46 AM		

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Widal Test (Slide Agglutination)

Salmonella typhi O (TO)	No Agglutination	Titre	< 1:80
Salmonella typhi H (TH)	No Agglutination	Titre	< 1:160
Salmonella paratyphi A(H)	No Agglutination	Titre	< 1:80
Salmonella Paratyphi B(H)	No Agglutination	Titre	< 1:80

Interpretation:

METHOD-(Slide Agglutination)

1. Titres >1:80 of "O" antigen & >1:160 of "H" antigen for Salmonella typhi and titres >1:80 of "H" antigen for Salmonella paratyphi A & B are reactive.
2. Rising titres in paired samples taken 7-10 days apart are more significant than a single test.
3. Reactive results indicates ongoing or recent infection by Salmonella spp. and the diagnosis should be confirmed by gold standard test such as Blood culture.
4. The reactivity will vary with stage of the disease with appearance in 1st week to increase in titres till end of 4th week post which it starts decreasing.
5. In TAB vaccinated patients, high titres of H antibody of $\geq 1:160$ to each of Salmonellae is observed. They tend to persist for many months and even years while O antibody shows lower titres and disappears within 6 months.
6. Antibiotic treatment during 1st week before the appearance of antibodies tend to suppress the immune response in the form of no or decreasing antibody levels.
7. False positive results/anamnestic response may be seen in patients with past enteric infection and during unrelated fevers like Malaria, Influenzae etc. in the form of transient rise in H antibody in Widal test.
8. False negative results may be due to processing of sample collected early in the course of disease (1st week) and immunosuppression.
9. Test conducted on serum.

Uses

- To diagnose infection due to Salmonella spp. (Enteric fever).
- To monitor the progression of disease.
- To assess the response to therapy (decreasing titres) in patients being treated for Enteric fever

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

K. Kokila

Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO :
Patient ID / UHID :		Sample Type : Serum
Referred BY : Self		Report Date : Feb 26, 2026, 08:29 PM.
Sample Collected : Feb 26, 2026, 11:46 AM		

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Dengue Ns1 Antigen Test, EIA

DENGUE NS1 ANTIGEN (Serum,EIA)	0.025	Ag Unit	Negative <1 Positive >=1
-----------------------------------	-------	---------	-----------------------------

Interpretation:

BIOLOGICAL REFERANCE INTERVAL

NEGATIVE <1

POSITIVE >=1

Note: As per regulation, specimen collecting Laboratory is responsible for reporting positive Dengue cases to Municipal corporation.

Indication: The Dengue (NS1) Antigen assay is a Enzyme linked immunoassay (EIA) for the detection of Dengue virus NS1 Antigen in human serum or plasma(heparin).

The serological detection of the highly specific dengue virus NS1antigen in patients with a dengue virus infection is possible at the onset of clinical symptoms in primary as well as secondary infections. Thus determination of Dengue (NS1) Antigen is an important supportive aid for diagnosis of acute dengue virus infections.

Clinical background: Dengue virus (serotypes Dn 1, Den 2, Den 3, Den 4) is a flavivirus with global distribution and is transmitted by mosquitoes (Aedes aegyptii, Aedes albopictus etc). It may cause Dengue fever, Dengue haemorrhagic fever or Dengue Shock syndrome.

Following the dengue infection, an incubation period of 3 to7 days, some infections maybe asymptomatic. Symptomatic patients develop fever with or without rash, severe musculoskeletal pain, headache, retro-orbital pain, petechiae etc. In most individuals there is resolution of illness without complications. In some individuals the Dengue fever may progress to Dengue haemorrhagic fever or Dengue Shock syndrome especially during repeat infection with a new Dengue Virus serotype.

Dengue virus antigen usually appears in blood within 24 hours of onset of symptoms to symptoms till 9 days post onset of symptoms.

Positive: The presence of Dengue nonstructural protein 1 (NS1) antigen is consistent with acute infection with dengue virus. The NS1 antigen is typically detectable within 1 to 2 days following infection and up to 9 days following symptom onset. NS1 antigen may also be detectable during secondary dengue virus infection, but for a shorter duration of time (1-4 days following symptom onset).

Negative: The absence of dengue NS1 antigen is suggestive of absence of acute phase of the infection. The NS1 antigen may be negative if specimen is collected too early such as immediately following dengue virus infection (<24-48 hours) or is collected following 9 to 10 days of symptoms. Results should always be interpreted in conjunction with clinical presentation and exposure history.

Limitations: Uncommonly, false positive Dengue NS1 antigen results may be seen in individuals with other flaviviruses west nile virus as well as Yellow fever. Negative NS1 antigen results may occur if the specimen was collected greater than 7 days following symptom onset. Serologic testing for the presence of IgM and IgG antibodies to Dengue Virus is recommended in such cases.

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

K. Kokila

Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO :
Patient ID / UHID :		Sample Type : Serum
Referred BY : Self		Report Date : Feb 26, 2026, 06:56 PM.
Sample Collected : Feb 26, 2026, 11:46 AM		

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Dengue IgM Antibodies, EIA

DENGUE FEVER ANTIBODY, IgM, SERUM SERUM, EIA	0.041	Units	Negative <1 Positive >=1
---	-------	-------	-----------------------------

Interpretation:

RESULT(Units)	REMARKS
Negative (<1)	No detectable IgM antibody. Result does not rule out Dengue infection. Additional sample to be tested after 7-14 days if infection is suspected.
Positive (>1)	IgM antibody detected. Suggestive of Primary / Secondary Dengue infection.

NOTE-

- The test should be used for detection of IgM antibodies of dengue in human serum/plasma.
- This is only a screening test and will only indicate the presence or absence of Dengue antibodies in the specimen. All reactive sample should be confirmed by confirmatory test. Therefore for a definitive diagnosis, the patients clinical history , symptomatology as well as serological data should be considered. The results should be reported only after complying with the above procedure.
- False positive results can be obtained due to cross reaction with Epstein-BARR virus, RA, Leptospira, Malaria, hepatitis-A, Influenza A & B, S.typhi Japanese encephalites, westnile virus diseased. This occurs in less then 1% of the sample tested.
- Immuno-depressive treatments presumably after the immune response to infection, inducing negative results in IgM in dengue patients.

Comments

Dengue viruses belong to the family Flaviviridae and have 4 subtypes (1-4). Dengue virus is transmitted by the mosquito Aedes aegypti and Aedes albopictus, widely distributed in Tropical and Subtropical areas of the world. Dengue is considered to be the most important arthropod borne viral disease due to the human morbidity and mortality it causes. The disease may be subclinical, self limiting, febrile or may progress to a severe form of Dengue hemorrhagic fever or Dengue shock syndrome.

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

K. Kokila

Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO :
Patient ID / UHID :		Sample Type : Serum
Referred BY : Self		Report Date : Feb 26, 2026, 06:56 PM.
Sample Collected : Feb 26, 2026, 11:46 AM		

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Dengue IgG Antibodies, EIA

DENGUE FEVER ANTIBODY, IgG, SERUM (Serum, EIA)	0.032	Units	Negative <1 Positive >=1
---	-------	-------	-----------------------------

Interpretation:

RESULT(Units)	REMARKS
Negative (<1)	No detectable IgG antibody. Result does not rule out Dengue infection. Additional sample to be tested after 7-14 days if infection is suspected.
Positive (>1)	IgG antibody detected. Suggestive of Primary / Secondary Dengue infection.

- NOTE-
- The test should be used for detection of IgG antibodies of dengue in human serum/plasma.
 - This is only a screening test and will only indicate the presence or absence of Dengue antibodies in the specimen. All reactive sample should be confirmed by confirmatory test. Therefore for a definitive diagnosis, the patients clinical history , symptomatology as well as serological data should be considered. The results should be reported only after complying with the above procedure.
 - False positive results can be obtained due to cross reaction with Epstein-BARR virus, RA, Rubella, Anti-nuclear antibody, Japanese encephalites, westnile virus diseased. This occurs in less then 1% of the sample tested.
 - Immuno-despressive treatments presumably after the immune response to infection, inducing negative results in IgG in dengue patients.

Comments

Dengue viruses belong to the family Flaviviridae and have 4 subtypes (1-4). Dengue virus is transmitted by the mosquito Aedes aegypti and Aedes albopictus, widely distributed in Tropical and Subtropical areas of the world. Dengue is considered to be the most important arthropod borne viral disease due to the human morbidity and mortality it causes. The disease may be subclinical, self limiting, febrile or may progress to a severe form of Dengue hemorrhagic fever or Dengue shock syndrome.

Chikungunya IgM, Rapid Card

Chikungunya IgM, Rapid Card Immunochromatographic Assay	Negative	Negative
--	----------	----------

Interpretation:

- Note:**
- It is a screening test
 - The Onsite Chikungunya IgM Rapid test is limited to the qualitative detection of IgM in human sera .
 - A negative result indicates absence of detectable IgM antibodies. However a negative result does not preclude the possibility of exposure or infection with CHIK.
 - Positive results in serum indicates ongoing or recent infection by Chikungunya virus causing rash, fever and severe joint pain (arthralgia). The test cannot be used to differentiate between primary and secondary infection.
 - Any reactive or non reactive report is recommended to be confirmed with additional test like PCR.
 - False positive results may be seen due to cross reactivity in infection with Epstein -BARR virus, Influenza A & B, Brucella and other mosquito borne diseases like Dengue and Zika Virus infections, in patients with high levels of heterophile antibodies and Rheumatoid factor.
 - False negative results may be due to processing of sample collected early in the course of disease and immunosuppression.
 - Test conducted on serum.

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

K. Kokila

Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO :
Patient ID / UHID :		Sample Type : Serum
Referred BY : Self		Report Date : Feb 26, 2026, 06:56 PM.
Sample Collected : Feb 26, 2026, 11:46 AM		

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Typhidot IgG & IgM, Rapid Card

TYPHI DOT/ SALMONELLA TYPHI IgG <i>Qualitative immunoassay, rapid card</i>	Negative	-	Negative
TYPHI DOT/ SALMONELLA TYPHI IgM <i>Qualitative immunoassay, rapid card</i>	Negative	-	Negative

Interpretation:

RESULTS	REMARKS
Positive	Indicates presence of IgM & IgG antibodies against Salmonella typhi.
Negative	Indicates absence of IgM & IgG antibodies against Salmonella spp.

- Note:**
1. Its positivity in serum indicates ongoing or recent infection by Salmonella typhi and the diagnosis should be confirmed by gold standard test such as Blood culture prior to start of antibiotics.
 2. IgM antibodies are typically detectable 5-7 days post symptom onset, peaking in 2nd week and frequently remain elevated for 2-4 months following infection.
 3. False positive results may be due to cross reactivity with other Salmonella spp., Dengue virus infection & in patients with high levels of Rheumatoid factor.
 4. False negative reaction may be due to processing of sample collected early in the course of disease, antibiotic treatment during 1st week and immunosuppression.
 5. Test conducted on serum.

Use
To diagnose infection due to Salmonella typhi (Enteric fever)

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.

K. Kokila

Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection
Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

Patient NAME		Report STATUS : Final Report
DOB/Age/Gender		Barcode NO
Patient ID / UHID		Sample Type : Spot Urine
Referred BY : Self		Report Date : Feb 26, 2026, 05:27 PM.
Sample Collected : Feb 26, 2026, 11:42 AM		

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Urine Routine and Microscopic Examination

Physical Examination			
Volume *	20	ml	-
Colour *	Pale yellow	-	Pale yellow
Transparency *	Clear	-	Clear
Deposit *	Absent	-	Absent

Chemical Examination			
Reaction (pH) <i>Double Indicator</i>	6.0	-	4.5 - 8.0
Specific Gravity <i>Ion Exchange</i>	1.025	-	1.010 - 1.030
Urine Glucose (sugar) <i>Oxidase / Peroxidase</i>	Negative	-	Negative
Urine Protein (Albumin) <i>Acid / Base Colour Exchange</i>	Negative	-	Negative
Urine Ketones (Acetone) <i>Legals Test</i>	Negative	-	Negative
Blood <i>Peroxidase Hemoglobin</i>	Negative	-	Negative
Leucocyte esterase <i>Enzymatic Reaction</i>	Negative	-	Negative
Bilirubin Urine <i>Coupling Reaction</i>	Negative	-	Negative
Nitrite <i>Griless Test</i>	Negative	-	Negative
Urobilinogen <i>Ehrlichs Test</i>	Normal	-	Normal


Microscopic Examination			
Pus Cells (WBCs) *	1-2	/hpf	0 - 5
Epithelial Cells *	1-2	/hpf	0 - 4
Red blood Cells *	Absent	/hpf	Absent
Crystals *	Absent	-	Absent
Cast *	Absent	-	Absent
Yeast Cells *	Absent	-	Absent
Amorphous deposits *	Absent	-	Absent
Bacteria *	Absent	-	Absent
Protozoa *	Absent	-	Absent

Interpretation:

URINALYSIS- Routine urine analysis assists in screening and diagnosis of various metabolic, urological, kidney and liver disorders.

Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria, dehydration, urinary tract infections and acute illness with fever

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection


Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

Patient NAME :		Report STATUS : Final Report
DOB/Age/Gender :		Barcode NO :
Patient ID / UHID :		Sample Type : Spot Urine
Referred BY : Self		Report Date : Feb 26, 2026, 05:27 PM.
Sample Collected : Feb 26, 2026, 11:42 AM		

Test Description	Value(s)	Unit(s)	Reference Range
<p>Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine. Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.</p> <p>Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine. Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise.</p> <p>Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders.</p> <p>Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys. Most common cause is bacterial urinary tract infection.</p> <p>Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration during infection increases with length of time the urine specimen is retained in bladder prior to collection.</p> <p>pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/ alkalosis or ingestion of certain type of food can affect the pH of urine.</p> <p>Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased specific gravity is seen in conditions like dehydration, glycosuria and proteinuria while decreased specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus.</p> <p>Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.</p> <p>Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in cases of haemolytic anaemia.</p>			

*** End Of Report ***

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Dr. K Kokila
MD, DNB(Path)
Consultant Pathologist

Booking Centre :- Home Collection

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., Plot No.402, Paneer Nagar, Mogappair Village, Ambattur TK Chennai - 600037

Terms and Conditions of Reporting

1. The presented findings in the Reports are intended solely for informational and interpretational purposes by the referring physician or other qualified medical professionals possessing a comprehensive understanding of reporting units, reference ranges, and technological limitations. The laboratory shall not be held liable for any interpretation or misinterpretation of the results, nor for any consequential or incidental damages arising from such interpretation.
2. It is to be presumed that the tests performed pertain to the specimen/sample attributed to the Customer's name or identification. It is presumed that the verification particulars have been cleared out by the customer or his/her representation at the point of generation of said specimen / sample. It is hereby clarified that the reports furnished are restricted solely to the given specimen only.
3. It is to be noted that variations in results may occur between different laboratories and over time, even for the same parameter for the same Customer. The assays are performed and conducted in accordance with standard procedures, and the reported outcomes are contingent on the specific individual assay methods and equipment(s) used, as well as the quality of the received specimen.
4. This report shall not be deemed valid or admissible for any medico-legal purposes.
5. The Customers assume full responsibility for apprising the Company of any factors that may impact the test finding. These factors, among others, includes dietary intake, alcohol, or medication / drug(s) consumption, or fasting. This list of factors is only representative and not exhaustive.

DISCLAIMER

This is a sample report provided for demonstration purposes only and does not represent an actual patient report. Test results, reference ranges, methodologies, instrumentation, and report formats may vary depending on the laboratory performing the test. The format and representation shown are indicative of reports generated by the National Reference Laboratory of Redcliffe Labs, Noida. This sample report should not be used for medical interpretation, diagnosis, or treatment decisions.