

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			
Test Description	Value(s)	Unit(s)	Reference Range

### Sexual Wellness Check up Panel

#### VDRL

VDRL RAPID CHROMATOGRAPHIC IMMUNOASSAY	NON REACTIVE	-	NON REACTIVE
---	--------------	---	--------------

#### Interpretation:

RESULTS	REMARKS
Reactive	Indicates presence of IgM & IgG antibodies against Treponemal Pallidum antigens
Non Reactive	Indicates absence of IgM & IgG antibodies against Treponemal Pallidum antigens

#### **Note**

1. Positive result indicates ongoing or recent infection and the diagnosis should be confirmed by specific Treponemal tests such as TPHA & FTA- AbS.
2. The reactivity will vary with Primary (60-86%), Secondary (99%) and Tertiary (98%) stage of Syphilis.
3. False positive results may be observed in patients of Malaria, Hepatitis, Mumps, Leprosy, Infectious Mononucleosis, Rheumatoid Arthritis and Collagen disease.
4. False negative reaction may be due to processing of sample collected early in the course of disease, immunosuppression and due to prozone effect.
5. Test conducted on serum.
6. It is a qualitative test.

#### **Uses**

To screen for presence of Syphilis infection.

*K. Kokila*

Dr. K Kokila  
MD, DNB(Path)  
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

### Chlamydia Trachomatis IgG Antibodies

Chlamydia Trachomatis IgG (Serum, EIA)	0.15	Ratio	<0.8
---	------	-------	------

**Interpretation:**


RESULT (RATIO)	REMARKS
<0.8	Negative
>=0.8 - <1.1	Equivocal
>=1.1	Positive

**Note**

1. Equivocal results do not rule out the possibility of Chlamydial infection. Retesting is recommended after 7 days.
2. Results must be correlated with clinical findings and other diagnostic investigations.

**Comments**

Chlamydia trachomatis is implicated in a wide variety of infections in humans. It is a common cause of Non-gonococcal urethritis and Cervicitis. In females it causes Pelvic Inflammatory disease, Salpingitis & Endometritis. In males it leads to Epididymitis & Reiter's syndrome. Lymphogranuloma venereum (LGV) is a sexually transmitted infection caused by Chlamydia trachomatis. It can also cause ophthalmologic infections like Trachoma and Inclusion Conjunctivitis.



Dr. Nitin Arora  
MBBS, MD (Microbiology)

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			
Test Description	Value(s)	Unit(s)	Reference Range

### Hepatitis C Antibody (HCV), Qualitative

Hepatitis C virus-Total Antibodies <i>CMIA</i>	0.1	S/CO	< 1.00 Nonreactive =>1.00 Reactive
---	-----	------	---------------------------------------

**Interpretation:**

Hepatitis C (HCV) is an RNA virus of Flavivirus group transmitted via blood transfusions, transplantation, injection drug users, accidental needle punctures in healthcare workers, dialysis patients and rarely from mother to infant. 10% of new cases show sexual transmission. As compared to HAV & HBV, chronic infection with HCV occurs in 85% of infected individuals. In high risk populations, the predictive value of Anti HCV for HCV infection is > 99% whereas in low risk populations it is only 25%.

**Note**

- 1.The Anti HCV Antibody assay is an in vitro diagnostic test for the qualitative detection of antibodies to hepatitis C virus (HCV) in human serum and plasma.
- 2.False positive results are seen in Autoimmune diseases, Rheumatoid factor, Hypergammaglobulinemia, Paraproteinemia, passive antibody transfer, Anti- idiotypes & Anti superoxide dismutase
- 3.False negative results are seen in early Acute infection, Immunosuppression & Immuno-incompetence.
- 4.HCV RNA PCR recommended in all Reactive results to differentiate between past and present infection.
- 5.It is a screening test and reactive samples need to confirmed by alternate method like viral load or PCR

### Herpes Simplex Virus (HSV) 1 + 2 IgG Antibody

Herpes simplex virus 1+2, IgG <i>EIA, SERUM</i>	0.1	INDEX	NEGATIVE: < 0.8 EQUIVOCAL: 0.8 and 1.2. POSITIVE: > 1.2
--	-----	-------	---

**Interpretation:**

RESULT IN INDEX	REMARKS
< 0.8	Negative
0.8 and 1.2.	Equivocal
≥ 1.2	Positive

The Herpes simplex virus (HSV) is a member of the Herpesviridae family, of which two types are known: type 1 (HSV-1) and type 2 (HSV-2) which present slight antigenic differences. HSV-1 causes chiefly oral-facial lesions, while HSV-2 is mainly responsible for genital lesions, but this distinction is not binding, both types occasionally causing infection in either anatomical site. HSV may also cause a form of ocular cheratitis, and lesions of the central nervous system. HSV can affect practically the whole population. The primary infection is often in a subclinical form and is rarely diagnosed. After a latency period of variable duration, reactivation may occur and viral replication may or may not give rise to clinical lesions. Infection contracted during birth is of particular interest, this being an important cause of morbidity and mortality. It is therefore important to determine the immunitary state of women during pregnancy in order to detect serum conversion.



Dr. Nitin Arora  
MBBS,MD (Microbiology)

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------


**Herpes Simplex Virus (HSV) 1 + 2 IgM Antibody**

Herpes simplex virus 1+2, IgM <i>EIA, SERUM</i>	0.4	Index	
--	-----	-------	--

**Interpretation:**

RESULT IN INDEX	REMARKS
< 0.9	Negative
0.9 and 1.1	Equivocal
> 1.1	Positive

The Herpes simplex virus (HSV) is a member of the Herpesviridae family, of which two types are known: type 1 (HSV-1) and type 2 (HSV-2) which present slight antigenic differences. HSV-1 causes chiefly oral-facial lesions, while HSV-2 is mainly responsible for genital lesions, but this distinction is not binding, both types occasionally causing infection in either anatomical site. HSV may also cause a form of ocular cheratitis, and lesions of the central nervous system. HSV can affect practically the whole population. The primary infection is often in a subclinical form and is rarely diagnosed. After a latency period of variable duration, reactivation may occur and viral replication may or may not give rise to clinical lesions. Infection contracted during birth is of particular interest, this being an important cause of morbidity and mortality. It is therefore important to determine the immunitary state of women during pregnancy in order to detect serum conversion. The assay of specific IgM is important for the diagnosis of neonatal infection and encephalitis caused by HSV. Moreover, the presence of specific IgM indicates viral activity in progress, although it is not possible to distinguish between primary infection and reactivation.



**Dr. Nitin Arora**  
MBBS,MD (Microbiology)

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

### Treponema Pallidum Hemagglutination Assay (TPHA)

Treponema Pallidum Haemagglutination (TPHA) LATEX AGGLUTINATION (QUALITATIVE)	Negative		Negative
--	----------	--	----------

**Interpretation:**

Test performed at 1:80 starting dilution

Result	Remarks
Positive	Indicate presence of specific Treponema pallidum antibodies in patient sera
Negative	Indicate absence of specific Treponema pallidum antibodies in patient sera

**Note:-**

1. A positive TPHA result cannot differentiate past or current T. pallidum infection thus quantitative RPR test is recommended for further evaluation of current infection or for patients on treatment.
2. A negative TPHA result generally indicates no current or past syphilis exposure. However, false negative results can be obtained during early -stage infections or window period and may require follow-up testing.
3. False positive TPHA results may be observed in infections with other treponemal species, leprosy, infectious mononucleosis, H. pylori infection, toxoplasmosis, and/or connective tissue disorders.
4. False negative TPHA results can occur in the early stages of syphilis or in immuno-compromised individuals (unable to mount immune response).

**Interpretation:-**


Syphilis is a chronic infection that progresses through stages of infection, primary to quaternary. The typical symptoms initially are sores known as chancres, then syphilitic rash followed by long periods of dormancy. Untreated infection may result in cardiovascular problems and neurosyphilis. The infection is caused by the Spirochaete Treponema pallidum, and is usually acquired by sexual contact, although the disease may be transmitted by transfusion of infected blood. Intrauterine infection also occurs. The organism has proved virtually impossible to culture in artificial media and diagnosis of the infection usually depends on the demonstration of antibodies in the blood, which appear soon after initial infection.

### Hepatitis B Surface Antigen (HBsAg), Qualitative

Hepatitis B surface antigen (HBs Ag) CMIA	0.3	S/CO	< 1.00 Nonreactive > 1.00 Reactive
--	-----	------	---------------------------------------

**Interpretation:**

Infection with HBV results in a wide spectrum of acute & chronic liver diseases & also is clearly linked with the development of hepatocellular carcinoma. HBV infection produces an array of unique antigens and antibodies which follow distinct and individual serological patterns. By monitoring these markers, it is possible not only to diagnose infection, but also to determine the stage of the disease and probable prognosis. HBsAg is the first marker to appear following infection, and is the best indirect indicator of potentially infectious sera. It is a screening test and reactive samples need to be confirmed by alternate method like viral load or PCR



Dr. Nitin Arora  
MBBS,MD (Microbiology)

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

**HIV 1&2 Antibody, Qualitative**

HIV I & II Qualitative & P24 COMBO CMIA	0.1	S/CO	<1.00 :Nonreactive >1.00 :Reactive
--	-----	------	---------------------------------------

**Interpretation:**  
The cutoff is 1.00 S/CO.


S/CO	Reference Ranges
< 1.00	Nonreactive
≥ 1.00	Reactive

**Final Interpretation**

Initial Interpretation	Results with Retes	Final Interpretation
Nonreactive	No retest required	Nonreactive. HIV p24 Ag and/or HIV-1/HIV-2 Ab not detected.
Reactive	If both retest results are < 1.00	Nonreactive. HIV p24 Ag and/or HIV-1/HIV-2 Ab not detected
	If one or both retest results are ≥ 1.00	Reactive. Presumptive evidence of HIV p24 Ag and/or HIV-1/ HIV-2 Ab; perform a supplemental assay

1. Non-Reactive result implies that no Anti HIV-I OR HIV-II Antibodies have been detected in the sample by this method. This means that either the patient has not exposed to HIV-I or HIV-II infection or the sample has been tested during the "WINDOW PHASE"(before the development of detectable levels of antibodies). Hence negative result does not exclude the possibility of exposure to or infection with HIV-I/HIV-II.
2. Immunoassay for the in vitro qualitative determination of HIV-1 p24 antigen and antibodies to HIV-1, including group O, and HIV-2 in human serum
3. It is a screening test and reactive samples need to confirmed by alternate method like western blot or PCR

\*\*\* End Of Report \*\*\*



Dr. Nitin Arora  
MBBS,MD (Microbiology)

# Terms and Conditions of Reporting

1. The presented findings in the Reports are intended solely for informational and interpretational purposes by the referring physician or other qualified medical professionals possessing a comprehensive understanding of reporting units, reference ranges, and technological limitations. The laboratory shall not be held liable for any interpretation or misinterpretation of the results, nor for any consequential or incidental damages arising from such interpretation.
2. It is to be presumed that the tests performed pertain to the specimen/sample attributed to the Customer's name or identification. It is presumed that the verification particulars have been cleared out by the customer or his/her representation at the point of generation of said specimen / sample. It is hereby clarified that the reports furnished are restricted solely to the given specimen only.
3. It is to be noted that variations in results may occur between different laboratories and over time, even for the same parameter for the same Customer. The assays are performed and conducted in accordance with standard procedures, and the reported outcomes are contingent on the specific individual assay methods and equipment(s) used, as well as the quality of the received specimen.
4. This report shall not be deemed valid or admissible for any medico-legal purposes.
5. The Customers assume full responsibility for apprising the Company of any factors that may impact the test finding. These factors, among others, includes dietary intake, alcohol, or medication / drug(s) consumption, or fasting. This list of factors is only representative and not exhaustive.

---

## DISCLAIMER

This is a sample report provided for demonstration purposes only and does not represent an actual patient report. Test results, reference ranges, methodologies, instrumentation, and report formats may vary depending on the laboratory performing the test. The format and representation shown are indicative of reports generated by the National Reference Laboratory of Redcliffe Labs, Noida. This sample report should not be used for medical interpretation, diagnosis, or treatment decisions.