

smart Health Report

An Insightful Health Analytics Report
for Easier Understanding

Prepared For



Name Gender

Patient ID Age

Your Health Summary

Understand Your Health At A Glance
Your Personalized Health Summary is Now Available.

Your Health at a Glance – A Personalized Journey

Report Sections

- 1 Body Summary**
 A visual snapshot of your overall health, simple and easy to understand
- 2 Quick Health Highlights**
 Your health scores and a single view of all abnormal results for quick attention
- 3 Lab Report Overview**
 Understand at a glance which tests are normal and which are abnormal
- 4 Comparative Health Insights**
 See what has improved and what needs attention
- 5 Personalized Health Advisory**
 Actionable insights and expert guidance tailored just for you
- 6 Doctor's Reference Report**
 Complete lab results with interpretations to share with your healthcare provider

How to Read This Report

This comprehensive health report provides detailed insights into your test results. Each section offers different perspectives on your health status, from visual summaries to detailed analysis and personalized recommendations.


Name _____ Gender _____
Patient ID _____ Age _____

Your Health Summary

Understand Your Health At A Glance
Your Personalized Health Summary is Now Available.

● All In Range ● Borderline ● Out Of Range

Health Summary



Note: This section offers a quick snapshot of selected parameters. For all parameters and detailed analysis with clinical interpretation, please refer to the following pages

Name Gender

Patient ID Age

Your Health Summary

Understand Your Health At A Glance
Your Personalized Health Summary is Now Available.




Quick Health Summary

Personal Insights - Health Score

63

Overall, most parameters are within normal ranges, indicating a generally balanced health status. The Diabetes and Anemia profiles may affect energy levels and overall well-being, so consider maintaining a balanced diet rich in fruits, vegetables, and whole grains, along with regular physical activity such as walking or yoga. Routine check-ups are advisable to monitor these areas, and consulting a healthcare professional can provide personalized guidance. Remember, small lifestyle changes can lead to meaningful improvements in your health and vitality.

Note - Higher scores tentatively indicate better health status



Summary of Key Health Indicators

Total Parameters Tested	Borderline Results	Out Of Range Results
96	13	11

Health Status by Body System

Profile	Total	Borderline	Out of Range	Key Results
Cardiac Profile	10	2	5	<ul style="list-style-type: none"> ● HsCRP (2.89) ● Triglycerides (288.1) ● HDL Cholesterol (30.5) ● VLDL (57.62) ● HDL : LDL ratio (0.39) ● Non - HDL Cholesterol (136.5) ● Total Cholesterol : HDL ratio (5.48)
Blood Disorder	16	5	2	<ul style="list-style-type: none"> ● Lymphocytes (49) ● Abs. Basophil Count (0) ● Abs. Lymphocyte Count (3.09) ● MPV (14.4) ● PDW (25.9) ● P-LCR (58.9) ● P-LCC (39)
Inflammation	1	0	1	<ul style="list-style-type: none"> ● ESR (27)

Profile	Total	Borderline	Out of Range	Key Results
Iron	4	1	1	<ul style="list-style-type: none"> ● % Saturation Transferrin (63.97) ● TIBC (245.1)
Vitamin Profile	2	0	1	<ul style="list-style-type: none"> ● Vitamin B12 (151)
Infectious Diseases	4	0	1	<ul style="list-style-type: none"> ● PCT (0.1)
Diabetes Monitoring	4	1	0	<ul style="list-style-type: none"> ● Blood Sugar (Fasting) (104.8)
Thyroid Profile	3	0	0	All In Range
Anemia Studies	9	1	0	<ul style="list-style-type: none"> ● Haemoglobin (11.2)
Liver Profile	14	1	0	<ul style="list-style-type: none"> ● Globulin (3.6)
Kidney Profile	12	2	0	<ul style="list-style-type: none"> ● Blood Urea Nitrogen (BUN) (9.72) ● Serum Creatinine (0.7)
Electrolytes	4	0	0	All In Range
Urinalysis	12	0	0	All In Range

Name Gender

Patient ID Age

Your Health Summary

Understand Your Health At A Glance
Your Personalized Health Summary is Now Available.

Report Summary ● In Range ● Borderline ● Out Of Range ● No color - Reference range not available

INFLAMMATION

Test Name	Result unit	Range
● ESR - Erythrocyte Sedimentation Rate	27 mm/hr	< 19

DIABETES MONITORING

Test Name	Result unit	Range
● Glycosylated Hemoglobin (HbA1c)	5 %	< 5.6
Estimated Average Glucose	96.8 md/dL	
● Glucose Fasting	104.8 mg/dL	70 - 100
Urine Glucose (sugar)	Negative	

IRON

Test Name	Result unit	Range
● Iron	156.8 µg/dL	50 - 170
● TIBC,(Total Iron Binding Capacity)	245.1 µg/dL	250 - 450
● UIBC	88.3 µg/dL	70 - 310
● Transferrin Saturation	63.97 %	14 - 50

CARDIAC PROFILE

Test Name	Result unit	Range
● HIGHLY SENSITIVE C-REACTIVE PROTEIN (hs-CRP)	2.89 mg/L	< 1
● Total Cholesterol	167 mg/dL	< 200
● Triglycerides	288.1 mg/dL	< 150
● HDL Cholesterol	30.5 mg/dL	40 - 80
● Non HDL Cholesterol	136.5 mg/dL	< 130
● LDL Cholesterol	78.88 mg/dL	30 - 100
● V.L.D.L Cholesterol	57.62 mg/dL	< 30
● ChoL/HDL Ratio	5.48 Ratio	3.5 - 5
● HDL/ LDL Ratio	0.39 Ratio	0.5 - 3
LDL/HDL Ratio	2.59 Ratio	

VITAMIN PROFILE

Test Name	Result unit	Range
● Vitamin - B12	151 pg/mL	200 - 1100
● Vitamin D 25 - Hydroxy	32.1 ng/mL	30 - 100

Name Gender

Patient ID Age

Your Health Summary

Understand Your Health At A Glance
Your Personalized Health Summary is Now Available.

Report Summary ● In Range ● Borderline ● Out Of Range ● No color - Reference range not available

THYROID PROFILE

Test Name	Result <small>unit</small>	Range
● Triiodothyronine (T3)	148 ng/dL	35 - 193
● Total Thyroxine (T4)	10.4 µg/dL	4.87 - 11.2
● Thyroid Stimulating Hormone (Ultrasensitive)	0.5 mIU/L	0.35 - 4.94

ANEMIA STUDIES

Test Name	Result <small>unit</small>	Range
● Hemoglobin	11.2 g/dL	12 - 15
● PCV	36.1 %	36 - 46
● MCV	96.8 fl	83 - 101
● MCH	29.9 pg	27 - 32
● MCHC	31.5 g/dL	31.5 - 34.5
● RDW (CV)	14.0 %	11.6 - 14
● RDW-SD	43.2 fl	35.1 - 43.9
Mentzer Index	26.16 %	
Red blood Cells	Absent /hpf	

BLOOD DISORDER

Test Name	Result <small>unit</small>	Range
● TLC	6.3 10 ³ /µL	4 - 10
● Neutrophils	44 %	40 - 80
● Lymphocytes	49 %	20 - 40
● Monocytes	4 %	2 - 10
● Eosinophils	3 %	1 - 6
● Basophils	0 %	< 2
● Neutrophils.	2.77 10 ³ /µL	2 - 7
● Lymphocytes.	3.09 10 ³ /µL	1 - 3
● Monocytes.	0.25 10 ³ /µL	0.2 - 1
● Eosinophils.	0.19 10 ³ /µL	0.02 - 0.5
● Basophils.	0 10 ³ /µL	0.02 - 0.5
● Platelet Count	151 10 ³ /µL	150 - 410
● Mean Platelet Volume (MPV)	14.4 fL	9.3 - 12.1
● PDW	25.9 fL	8.3 - 25
● P-LCR	58.9 %	18 - 50
● P-LCC	39 10 ⁹ /L	44 - 140

Name Gender

Patient ID Age

Your Health Summary

Understand Your Health At A Glance
Your Personalized Health Summary is Now Available.

Report Summary ● In Range ● Borderline ● Out Of Range ● No color - Reference range not available

INFECTIOUS DISEASES

Test Name	Result <small>unit</small>	Range
● PCT	0.1 %	0.17 - 0.32
Deposit	Absent	
Yeast Cells	Absent	
Protozoa	Absent	

LIVER PROFILE

Test Name	Result <small>unit</small>	Range
● Bilirubin Total	0.8 mg/dL	0.2 - 1.2
● Bilirubin Direct	0.3 mg/dL	< 0.5
● Bilirubin Indirect	0.5 mg/dL	0.1 - 1
● SGOT/AST	24.8 U/L	5 - 34
● SGPT/ALT	27 U/L	< 55
SGOT/SGPT Ratio	0.92 %	
● Alkaline Phosphatase	91 U/L	40 - 150
● Total Protein	8.3 g/dL	6.4 - 8.3
● Albumin	4.7 gm/dL	3.8 - 5
● Globulin	3.6 g/dL	2.3 - 3.5
● Albumin :Globulin Ratio	1.31	1 - 2.1
● Gamma Glutamyl Transferase (GGT)	14.5 U/L	12 - 64
Bilirubin Urine	Negative	
Urobilinogen	Normal	

KIDNEY PROFILE

Test Name	Result <small>unit</small>	Range
● Blood Urea	20.8 mg/dL	18 - 55
● Bun	9.72 mg/dL	9.8 - 20.1
● Creatinine	0.7 mg/dL	0.72 - 1.25
eGFR (CKD-EPI)	105.28 ml/min/1.73 sq m	
● Bun/Creatinine Ratio	13.89	12 - 20
● Urea / Creatinine Ratio	29.71	25.68 - 42.8
● Uric Acid	5.6 mg/dL	2.6 - 6
● Calcium Serum	9.8 mg/dL	8.4 - 10.2
Urine Protein (Albumin)	Negative	
Blood	Negative	
Crystals	Absent	
Cast	Absent	

Name Gender

Patient ID Age

Your Health Summary

Understand Your Health At A Glance
Your Personalized Health Summary is Now Available.

Report Summary ● In Range ● Borderline ● Out Of Range ● No color - Reference range not available

ELECTROLYTE PROFILE

Test Name	Result <small>unit</small>	Range
● Phosphorus	3.9 mg/dL	2.3 - 4.7
● Sodium	137.6 mmol/L	136 - 145
● Potassium	4.2 mmol/L	3.5 - 5.1
● Chloride	106.5 mmol/L	98 - 107

URINALYSIS

Test Name	Result <small>unit</small>	Range
Volume	20 ml	
Colour	Pale yellow	
Transparency	Clear	
● Reaction (pH)	6.0	4.5 - 8
● Specific Gravity	1.010	1.01 - 1.03
Urine Ketones (Acetone)	Negative	
Leucocyte esterase	Negative	
Nitrite	Negative	
Pus Cells (WBCs)	1-2 /hpf	
Epithelial Cells	1-2 /hpf	
Amorphous deposits	Absent	
Bacteria	Absent	

Name Gender

Patient ID Age

Your Health Summary

Understand Your Health At A Glance
Your Personalized Health Summary is Now Available.

Comparative Health Summary

● In Range ● Borderline ● Out Of Range

Personal Health Score Change

Your health score is **63/100** (09-04-2026)

Summary of Key Improvements / Declines	Outcome
Total parameters improved	5 of 65 parameters tested earlier
<ul style="list-style-type: none"> Vitamin D 25 - Hydroxy Thyroid Stimulating Hormone (Ultrasensitive) PCV MCHC Albumin 	
New Out of range parameters detected	9 new issues
<ul style="list-style-type: none"> HIGHLY SENSITIVE C-REACTIVE PROTEIN (hs-CRP) Lymphocytes PDW Glucose Fasting Globulin Bun Non HDL Cholesterol Chol/HDL Ratio HDL/ LDL Ratio 	

Parameter-Wise Comparison

Parameter	Current <small>09-04-2026</small>	Previous	Range	Value Change	Trend
ESR - Erythrocyte Sedimentation Rate	● 27	● 26 <small>27-07-2025</small>	0-19 mm/hr	+1	Still out of range
HIGHLY SENSITIVE C-REACTIVE PROTEIN (hs-CRP)	● 2.89	● 0.4 <small>27-07-2025</small>	0-1 mg/L	+2.5	Need Attention
Vitamin - B12	● 151	● 109.9 <small>27-07-2025</small>	200-1100 pg/mL	+41.1	Still out of range
Vitamin D 25 - Hydroxy	● 32.1	● 19.79 <small>27-07-2025</small>	30-100 ng/mL	+12.3	Improved
Thyroid Stimulating Hormone (Ultrasensitive)	● 0.5	● 31.34 <small>27-07-2025</small>	0.35-4.94 mIU/L	-30.8	Improved
Hemoglobin	● 11.2	● 9.9 <small>27-07-2025</small>	12.0-15.0 g/dL	+1.3	Still out of range
PCV	● 36.1	● 32.6 <small>27-07-2025</small>	36-46 %	+3.5	Improved
MCHC	● 31.5	● 30.3 <small>27-07-2025</small>	31.5-34.5 g/dL	+1.2	Improved

Parameter	Current 09-04-2026	Previous	Range	Value Change	Trend
Lymphocytes	● 49	● 37 27-07-2025	20-40 %	+12	Need Attention
Lymphocytes.	● 3.09	● 3.59 27-07-2025	1-3 10 ³ /μl	-0.5	Still out of range
Basophils.	● 0	● 0 27-07-2025	0.02-0.5 10 ³ /μl	0	Still out of range
Mean Platelet Volume (MPV)	● 14.4	● 14 27-07-2025	9.3-12.1 fL	+0.4	Still out of range
PCT	● 0.1	● 0.1 27-07-2025	0.17-0.32 %	0	Still out of range
PDW	● 25.9	● 24.5 27-07-2025	8.3-25 fL	+1.4	Need Attention
P-LCR	● 58.9	● 54.9 27-07-2025	18-50 %	+4	Still out of range
P-LCC	● 39	● 39 27-07-2025	44-140 10 ⁹ /L	0	Still out of range
Glucose Fasting	● 104.8	● 98.3 27-07-2025	70-100 mg/dL	+6.5	Need Attention
Albumin	● 4.7	● 3.7 27-07-2025	3.8-5 gm/dL	+1	Improved
Globulin	● 3.6	● 3.1 27-07-2025	2.3-3.5 g/dL	+0.5	Need Attention
Bun	● 9.72	● 10.84 27-07-2025	9.8-20.1 mg/dL	-1.1	Need Attention
Creatinine	● 0.7	● 0.6 27-07-2025	0.72-1.25 mg/dL	+0.1	Still out of range
Triglycerides	● 288.1	● 371.9 27-07-2025	0-150 mg/dL	-83.8	Still out of range
HDL Cholesterol	● 30.5	● 30.8 27-07-2025	40-80 mg/dL	-0.3	Still out of range
Non HDL Cholesterol	● 136.5	● 117.2 27-07-2025	0-130 mg/dL	+19.3	Need Attention
V.L.D.L Cholesterol	● 57.62	● 74.38 27-07-2025	0-30 mg/dL	-16.8	Still out of range
Chol/HDL Ratio	● 5.48	● 4.81 27-07-2025	3.5-5 Ratio	+0.7	Need Attention
HDL/ LDL Ratio	● 0.39	● 0.72 27-07-2025	0.5-3.0 Ratio	-0.3	Need Attention

Name

Gender

Patient ID

Age
--

 **Your Health Summary**

Understand Your Health At A Glance
Your Personalized Health Summary is Now Available.

Health Advisory

● In Range ● Borderline (BL) ● Out Of Range




Inflammation

Inflammation is the body's immune system's response to an injury, surgery, or irritation. This natural defense process acts by removing injurious stimuli and initiating the healing process. Inflammation can be chronic (such as arthritis) or acute (like in case of trauma).

ESR - Erythrocyte Sedimentation Rate: 27 mm/hr

● OUT OF RANGE






Diabetes

This panel is used to check how much glucose/sugar there is in your blood. Too much blood glucose might indicate diabetes.


Glycosylated Hemoglobin (HbA1c): 5%


● IN RANGE



Glucose Fasting: 104.8 mg/dL

● BORDERLINE






Cardiac Profile

Most people believe they are safe from heart diseases, but in reality, heart diseases are the leading cause of death in the world. There are many different forms of heart disease. Narrowing or blockage of the coronary arteries is the most common cause of heart disease, which are the vessels that supply blood to the heart. This is called coronary artery disease and it occurs slowly over time. It is the main cause of heart attacks.


HIGHLY SENSITIVE C-REACTIVE PROTEIN (hs-CRP): 2.89 mg/L

● OUT OF RANGE




Total Cholesterol: 167 mg/dL

● IN RANGE




Triglycerides: 288.1 mg/dL

● OUT OF RANGE




HDL Cholesterol: 30.5 mg/dL

● OUT OF RANGE



LDL Cholesterol: 78.88 mg/dL

● IN RANGE






Vitamins Profile

Vitamins are considered essential nutrients because they perform hundreds of roles in your body. They help maintain bones, heal wounds, and strengthen your immune system. They also convert food into energy, and repair cellular damage


Vitamin - B12: 151 pg/mL

● OUT OF RANGE



Vitamin D 25 - Hydroxy: 32.1 ng/mL

● IN RANGE






Thyroid

This panel is used to check the imbalance in your thyroid gland. A healthy thyroid gland is very important for metabolism, controlling body temperature, regulation of mood, muscle strength and regulation of body weight


Triiodothyronine (T3): 148_{ng/dL}

● IN RANGE




Total Thyroxine (T4): 10.4_{µg/dL}

● IN RANGE



Thyroid Stimulating Hormone (Ultrasensitive): 0.5_{mIU/L}

● IN RANGE






Anemia Profile

Anemia is the condition where your body has less RBCs (red blood cells) or the RBCs don't have enough haemoglobin. Haemoglobin is the protein present in RBCs that help carry oxygen to your body's tissues.

Hemoglobin: **11.2** g/dL

● BORDERLINE




Liver Profile


One of the main functions of your liver is to make proteins that are secreted in your blood. It also makes enzymes which convert food into energy, and processes old muscles and cells. When your liver is damaged, enzymes leak into your blood and appear in the blood test

Bilirubin Total: **0.8** mg/dL


● IN RANGE




Alkaline Phosphatase: 91 U/L ● IN RANGE



Albumin: 4.7 gm/dL ● IN RANGE




Gamma Glutamyl Transferase (GGT): 14.5 U/L ● IN RANGE




Enzymes

Enzymes found in your liver are responsible for various processes that maintain body functions. These enzymes are leaked into your blood when your liver suffers dysfunction.

SGOT/AST: 24.8 U/L ● IN RANGE



SGPT/ALT: 27 U/L ● IN RANGE






Kidney Profile

This panel is used to check healthy functioning of your kidneys. Kidneys filter blood in your body to remove waste products - these waste products are produced when breakdown of proteins (present in food, muscles and other cells) occurs in the body to generate energy


Bun: 9.72 mg/dL

● BORDERLINE




Creatinine: 0.7 mg/dL

● BORDERLINE




Uric Acid: 5.6 mg/dL

● IN RANGE



Calcium Serum: 9.8 mg/dL

● IN RANGE



Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			
Test Description	Value(s)	Unit(s)	Reference Range

Winter Special offer with Free Cardiac Screening

Complete Blood Count (CBC)

RBC Parameters			
Hemoglobin <i>colorimetric</i>	11.2 L*	g/dL	12.0 - 15.0
RBC Count <i>Electrical impedance</i>	3.7 L*	10 ⁶ /μl	3.8 - 4.8
PCV <i>Calculated</i>	36.1	%	36 - 46
MCV <i>Calculated</i>	96.8	fl	83 - 101
MCH <i>Calculated</i>	29.9	pg	27 - 32
MCHC <i>Calculated</i>	31.5	g/dL	31.5 - 34.5
RDW (CV) * <i>Calculated</i>	14.0	%	11.6 - 14.0
RDW-SD * <i>Calculated</i>	43.2	fl	35.1 - 43.9
WBC Parameters			
TLC <i>Electrical impedance and microscopy</i>	6.3	10 ³ /μl	4 - 10
Differential Leucocyte Count			
Neutrophils	44	%	40-80
Lymphocytes	49 H*	%	20-40
Monocytes	4	%	2-10
Eosinophils	3	%	1-6
Basophils	0	%	<2
Absolute Leukocyte Counts <i>Calculated</i>			
Neutrophils.	2.77	10 ³ /μl	2 - 7
Lymphocytes.	3.09 H*	10 ³ /μl	1 - 3
Monocytes.	0.25	10 ³ /μl	0.2 - 1.0
Eosinophils.	0.19	10 ³ /μl	0.02 - 0.5
Basophils.	0	10 ³ /μl	0.02 - 0.5
Platelet Parameters			
Platelet Count <i>Electrical impedance and microscopy</i>	151	10 ³ /μl	150 - 410
Mean Platelet Volume (MPV) * <i>Calculated</i>	14.4 H*	fL	9.3 - 12.1
PCT * <i>Calculated</i>	0.1 L*	%	0.17 - 0.32

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)



Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
PDW * <i>Calculated</i>	25.9 H*	fL	8.3 - 25.0
P-LCR * <i>Calculated</i>	58.9 H*	%	18 - 50
P-LCC * <i>Calculated</i>	39 L*	10 ⁹ /L	44 - 140
Mentzer Index * <i>Calculated</i>	26.16	%	-

Interpretation:

CBC provides information about red cells, white cells and platelets. Results are useful in the diagnosis of anemia, infections, leukemias, clotting disorders and many other medical conditions.

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)



Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Erythrocyte Sedimentation Rate (ESR)

ESR - Erythrocyte Sedimentation Rate <i>MODIFIED WESTERGREN</i>	27 H*	mm/hr	0 - 19
--	--------------	-------	--------

Interpretation:
 ESR is also known as Erythrocyte Sedimentation Rate. An ESR test is used to assess inflammation in the body. Many conditions can cause an abnormal ESR, so an ESR test is typically used with other tests to diagnose and monitor different diseases. An elevated ESR may occur in inflammatory conditions including infection, rheumatoid arthritis ,systemic vasculitis, anemia, multiple myeloma , etc. Low levels are typically seen in congestive heart failure, polycythemia ,sickle cell anemia, hypo fibrinogenemia , etc.

Reference- Dacie and lewis practical hematology

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)


Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

HbA1C (Glycosylated Haemoglobin)

Glycosylated Hemoglobin (HbA1c) <i>HPLC</i>	5	%	< 5.7
Estimated Average Glucose	96.8	md/dL	Refer Table Below

Interpretation:

Interpretation For HbA1c% As per American Diabetes Association (ADA)

Reference Group	HbA1c in %
Non diabetic adults >=18 years	<5.7
At risk (Prediabetes)	5.7 - 6.4
Diagnosing Diabetes	>= 6.5
Therapeutic goals for glycemic control	Age > 19 years Goal of therapy: < 7.0 Age < 19 years Goal of therapy: <7.5

Note:

- Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.
- Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments :

HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations ADA criteria for correlation between HbA1c & Mean plasma glucose levels.

HbA1c(%)	Mean Plasma Glucose (mg/dL)	HbA1c(%)	Mean Plasma Glucose (mg/dL)
6	126	12	298
8	183	14	355
10	240	16	413

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Blood Sugar Fasting

Glucose Fasting <i>Hexokinase</i>	104.8 H*	mg/dL	70 - 100
--------------------------------------	-----------------	-------	----------

Interpretation:

Status	Fasting plasma glucose in mg/dL
Normal	70 - 100
Impaired fasting glucose	101 - 125
Diabetes	≥126

Reference : American Diabetes Association

Comment :

Blood glucose determinations are commonly used as an aid in the diagnosis and treatment of diabetes. Elevated glucose levels (hyperglycemia) may also occur with pancreatic neoplasm, hyperthyroidism, and adrenal cortical hyper function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy insulinoma, or various liver diseases.

Note

- 1.The diagnosis of Diabetes requires a fasting plasma glucose of > or = 126 mg/dL or a random / 2 hour plasma glucose value of > or = 200 mg/dL with symptoms of diabetes mellitus.
- 2.Very high glucose levels (>450 mg/dL in adults) may result in Diabetic Ketoacidosis.

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)

P. Datta
Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Liver Function Test (LFT)

Bilirubin Total <i>Photometric</i>	0.8	mg/dL	0.2 - 1.2
Bilirubin Direct * <i>Diazo Reaction</i>	0.3	mg/dL	0.0 - 0.5
Bilirubin Indirect * <i>Calculation (T Bil - D Bil)</i>	0.5	mg/dL	0.1 - 1.0
SGOT/AST <i>IFCC without P5P</i>	24.8	U/L	5 - 34
SGPT/ALT <i>IFCC without P5P</i>	27	U/L	0 to 55
SGOT/SGPT Ratio *	0.92	-	-
Alkaline Phosphatase <i>p-nitrophenyl Phosphate, AMP buffer</i>	91	U/L	40 - 150
Total Protein <i>Biuret</i>	8.3	g/dL	6.4 - 8.3
Albumin <i>BCG</i>	4.7	gm/dL	3.8 - 5.0
Globulin * <i>Calculation (T.P - Albumin)</i>	3.6 H*	g/dL	2.3 - 3.5
Albumin :Globulin Ratio * <i>Calculation (Albumin/Globulin)</i>	1.31	-	1.0 - 2.1
Gamma Glutamyl Transferase (GGT) * <i>ENZYMATIC</i>	14.5	U/L	12 - 64

Interpretation:

The liver filters blood, metabolizes nutrients, detoxifies harmful substances, and produces blood clotting proteins. Liver cells contain enzymes that facilitate these functions. When cells are damaged, enzymes leak into the blood, detectable through blood tests.

Key enzymes tested:

- 1. AST (SGOT):** may indicate tissue injury / damage in muscles or liver.
- 2. ALT (SGPT):** Primarily in the liver. Elevated ALT and AST suggest liver damage.
- 3. Alkaline Phosphatase & GGT:** Linked to bile production and flow. Elevated levels may indicate bile flow issues related to the liver, gallbladder, or bile ducts.

Blood proteins, **albumin and globulin**, are essential for growth, development, and health.

- 1. Low protein:** May indicate bleeding, liver disorders, malnutrition, or agammaglobulinemia.
- 2. High protein (Hyperproteinemia):** Often due to dehydration or increased protein production.
- 3. Low albumin:** Caused by poor diet, kidney, or liver disease.
- 4. High albumin:** Usually due to severe dehydration.

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)



Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Kidney Function Test (KFT)

Blood Urea <i>Modified Jaffe</i>	20.8	mg/dL	18 - 55
Bun * <i>Urease</i>	9.72 L*	mg/dL	9.8 - 20.1
Creatinine <i>Modified Jaffe</i>	0.7 L*	mg/dL	0.72 - 1.25
eGFR (CKD-EPI) *	105.28	ml/min/1.73 sq m	Normal Or High: >= 90 Mild Or Decrease: 60-89 Mild To Moderate Decrease: 45-59 Mild To Severe Decrease: 30-44 Severe Decrease: 15-29 Kidney Failure: < 15
Bun/Creatinine Ratio * <i>Calculated</i>	13.89		12 - 20
Urea / Creatinine Ratio * <i>Calculated</i>	29.71		25.68 - 42.8
Uric Acid <i>Uricase</i>	5.6	mg/dL	2.6 - 6.0
Calcium Serum <i>O-Cresolphthalein Complex</i>	9.8	mg/dL	8.4 - 10.2
Phosphorus <i>Colorimetric - Phosphomolybdate Formation</i>	3.9	mg/dL	2.3 - 4.7
Sodium <i>Direct ISE</i>	137.6	mmol/L	136 - 145
Potassium <i>Direct ISE</i>	4.2	mmol/L	3.5 - 5.1
Chloride <i>Direct ISE</i>	106.5	mmol/L	98 - 107

Interpretation:

Kidney function tests is a collective term for a variety of individual tests and procedures that can be done to evaluate how well the kidneys are functioning. Many conditions can affect the ability of the kidneys to carry out their vital functions. Some lead to a rapid (acute) decline in kidney function others lead to a gradual (chronic) decline in function. Both result in a buildup of toxic waste substances done on urine samples, as well as on blood samples. A number of symptoms may indicate a problem with your kidneys. These include : high blood pressure, blood in urine, frequent urges to urinate, difficulty beginning urination, painful urination, swelling in the hands and feet due to a buildup of fluids in the body. A single symptom may not mean something serious. However, when occurring simultaneously, these symptoms suggest that your kidneys are not working properly. Kidney function tests can help determine the reason. Ionized calcium this test if you have signs of kidney or parathyroid disease. The test may also be done to monitor progress and treatment of these diseases."eGFR test is applicable for patients aged 18 years or more."

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)



Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Lipid Profile

Total Cholesterol <i>Enzymatic - Cholesterol Oxidase</i>	167	mg/dL	<200
Triglycerides <i>Colorimetric - Lip/Glycerol Kinase</i>	288.1 H*	mg/dL	<150
HDL Cholesterol <i>Accelerator Selective Detergent</i>	30.5 L*	mg/dL	>40
Non HDL Cholesterol * <i>Calculated</i>	136.5 H*	mg/dL	<130
LDL Cholesterol * <i>Calculated</i>	78.88	mg/dL	<100
V.L.D.L Cholesterol * <i>Calculated</i>	57.62 H*	mg/dL	< 30
Chol/HDL Ratio * <i>Calculated</i>	5.48 H*	Ratio	3.5 - 5.0
HDL/ LDL Ratio * <i>Calculated</i>	0.39 L*	Ratio	0.5 - 3.0
LDL/HDL Ratio * <i>Calculated</i>	2.59	Ratio	-

Interpretation:

Lipid level assessments must be made following 9 to 12 hours of fasting, otherwise assay results might lead to erroneous interpretation. NCEP recommends of 3 different samples to be drawn at intervals of 1 week for harmonizing biological variables that might be encountered in single assays.

National Lipid Association Recommendations (NLA-2014)	Total Cholesterol (mg/dL)	Triglyceride (mg/dL)	LDL Cholesterol (mg/dL)	Non HDL Cholesterol (mg/dL)
Optimal	<200	<150	<100	<130
Above Optimal			100-129	130 - 159
Borderline High	200-239	150-199	130-159	160 - 189
High	>=240	200-499	160-189	190 - 219
Very High	-	>=500	>=190	>=220

HDL Cholesterol	
Low	High
<40	>=60

Risk Stratification for ASCVD (Atherosclerotic Cardiovascular Disease) by Lipid Association of India.

Risk Category	A. CAD with > 1 feature of high risk group
Extreme risk group	B. CAD with >1 feature of very high risk group of recurrent ACS (within 1 year) despite LDL-C <or = 50 mg/dl or poly vascular disease
Very High Risk	1.Established ASCVD 2.Diabetes with 2 major risk factors of evidence of end organ damage 3. Familial Homozygous Hypercholesterolemia
	1. Three major ASCVD risk factors 2. Diabetes with 1 major risk factor or no evidence

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)


Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
High Risk	of end organ damage 3. CHD stage 3B or 4. 4 LDL >190 mg/dl 5. Extreme of a single risk factor 6. Coronary Artery Calcium - CAC > 300 AU 7. Lipoprotein a >= 50 mg/dl 8. Non stenotic carotid plaque		
Moderate Risk	2 major ASCVD risk factors		
Low Risk	0-1 major ASCVD risk factors		
Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors			
1. Age >=45 years in Males & >= 55 years in Females	3. Current Cigarette smoking or tobacco use		
2. Family history of premature ASCVD	4. High blood pressure		
5. Low HDL			

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by Lipid Association of India in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group Category A	<50 (Optional goal <OR = 30)	<80 (Optional goal <OR = 60)	>OR = 50	>OR = 80
Extreme Risk Group Category B	>OR = 30	>OR = 60	> 30	> 60
Very High Risk	<50	<80	>OR = 50	>OR = 80
High Risk	<70	<100	>OR = 70	>OR = 100
Moderate Risk	<100	<130	>OR = 100	>OR = 130
Low Risk	<100	<130	>OR = 130*	>OR = 160

* After an adequate non-pharmacological intervention for at least 3 months.

References : Management of Dyslipidaemia for the Prevention of Stroke : Clinical practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology,2022,20,134-155.

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)


Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Iron Studies

Iron <i>Ferene</i>	156.8	µg/dL	50 - 170
TIBC,(Total Iron Binding Capacity) <i>Calculated</i>	245.1 L*	µg/dL	250 - 450
UIBC <i>Ferene</i>	88.3	µg/dL	70 - 310
Transferrin Saturation <i>Method :Derived from IRON and TIBC values</i>	63.97 H*	%	14 - 50

Interpretation:

Increased levels due to iron ingestion or ineffective erythropoiesis. Decreased levels due to infection, inflammation, malignancy, menstruation and Fe deficiency. Needs to be taken into consideration with TIBC. Transferrin Saturation:- Low level Transferrin Saturation can indicate iron deficiency, erythropoiesis, infection, or inflammation. High level Transferrin Saturation can indicate recent ingestion of dietary iron, ineffective erythropoiesis, haemochromatosis or liver disease. High TIBC, UIBC, or transferrin usually indicates iron deficiency, but they are also increased in pregnancy and with the use of oral contraceptives. Low TIBC, UIBC, or transferrin may occur if someone has: Hemochromatosis, Certain types of anemia due to accumulated iron, Malnutrition, kidney disease that causes a loss of protein in urine.

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)



Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

High Sensitivity C-Reactive Protein (Hs-CRP)

HIGHLY SENSITIVE C-REACTIVE PROTEIN (hs-CRP) <i>immunoturbidimetric</i>	2.89 H*	mg/L	< 1.00
--	----------------	------	--------

Interpretation:

Cardio CRP In mg/L	Cardiovascular Risk
<1	Low
1-3	Average
3-10	High
>10	Persistent elevation may represent Non cardiovascular inflammation

Note: To assess vascular risk, it is recommended to test hsCRP levels 2 or more weeks apart and calculate the average

Comments:

High sensitivity C Reactive Protein (hsCRP) significantly improves cardiovascular risk assessment as it is a strongest predictor of future coronary events. It reveals the risk of future Myocardial infarction and Stroke among healthy men and women, independent of traditional risk factors. It identifies patients at risk of first Myocardial infarction even with low to moderate lipid levels. The risk of recurrent cardiovascular events also correlates well with hsCRP levels. It is a powerful independent risk determinant in the prediction of incident Diabetes.

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)


Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Vitamin B12 / Cyanocobalamin

Vitamin - B12 <i>CLIA</i>	151 L*	pg/mL	200 - 1100
------------------------------	---------------	-------	------------

Interpretation:

Low Values are a sign of a vitamin B12 deficiency. People with this deficiency are likely to have or develop symptoms.

Causes of vitamin B12 deficiency include: Not enough vitamin B12 in diet (rare except with a strict vegetarian diet), Diseases that cause malabsorption (for example, celiac disease and Crohn's disease), Lack of intrinsic factor, Above normal heat production (for example, with hyperthyroidism), Pregnancy. Increased vitamin B12 levels are uncommon. Usually excess vitamin B12 is removed in the urine. Conditions that can increase B12 levels include: Liver disease (such as cirrhosis or hepatitis), Myeloproliferative disorders (for example, polycythemia vera and chronic myelocytic leukemia).

Vitamin B12: Low Levels can cause malabsorption, Lack of intrinsic factor, Above normal heat production (for example, with hyperthyroidism), Pregnancy. High Level Liver disease, Myeloproliferative disorders (for example, polycythemia vera and chronic myelocytic leukemia).

1. Out of 140 healthy indian population, 91% of Vitamin B 12 concentrations was at lower level: 59.00 pg/ml and upper level: 700.00 pg/ml

"Patients on Biotin supplement may have interference in some immunoassays. Ref: Arch Pathol Lab Med—Vol 141, November 2017. With individuals taking high dose Biotin (more than 5 mg per day) supplements, at least 8-hour wait time before blood draw is recommended."

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)



Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Vitamin D 25 Hydroxy

Vitamin D 25 - Hydroxy <i>CLIA</i>	32.1	ng/mL	Deficiency: <10 Insufficiency: 10 - 29 Sufficiency: 30 - 100 Toxicity: >100
---------------------------------------	------	-------	--

Interpretation:

25-Hydroxy vitamin D represents the main body reservoir and transport form. Mild to moderate deficiency is associated with Osteoporosis / Secondary Hyperparathyroidism while severe deficiency causes Rickets in children and Osteomalacia in adults. Prevalence of Vitamin D deficiency is approximately >50% specially in the elderly. This assay is useful for diagnosis of vitamin D deficiency and Hypervitaminosis D. It is also used for differential diagnosis of causes of Rickets & Osteomalacia and for monitoring Vitamin D replacement therapy.

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Thyroid Profile Total

Triiodothyronine (T3) <i>CLIA</i>	148	ng/dL	35 - 193
Total Thyroxine (T4) <i>CLIA</i>	10.4	µg/dL	4.87 - 11.2
Thyroid Stimulating Hormone (Ultrasensitive) <i>CMIA</i>	0.5	mIU/L	0.35 - 4.94

Interpretation:

Pregnancy	Reference Range TSH
1st Trimester	0.1 - 2.5
2nd Trimester	0.2 - 3.0
3rd Trimester	0.3 - 3.0

Clinical Use:

1. Diagnose Hypothyroidism & Hyperthyroidism
2. Monitor T4 therapy
3. Measure subnormal TSH levels

Increased TSH: Primary hypothyroidism, Subclinical hypothyroidism, TSH-dependent hyperthyroidism, Thyroid hormone resistance

Decreased TSH: Graves' disease, Autonomous thyroid hormone secretion, TSH deficiency

Thyroid malfunction (hyper or hypo) affects T3 & T4 levels. Pituitary or hypothalamic issues also influence thyroid activity.

1. **Primary Hypothyroidism:** High TSH levels.
2. **Secondary/Tertiary Hypothyroidism:** Low TSH levels.
3. **Euthyroid Sick Syndrome:** Abnormal thyroid test results due to non-thyroidal illnesses (NTI).

TBG levels are stable in healthy individuals but may be altered by pregnancy, estrogens, androgens, steroids, or glucocorticoids, causing inaccurate T3 & T4 readings.

TSH	T4	T3	Interpretation
High	Normal	Normal	Mild (subclinical) hypothyroidism
High	Low	Low Or Normal	Hypothyroidism
Low	Normal	Normal	Mild (subclinical) hyperthyroidism
Low	High Or Normal	High Or Normal	Hyperthyroidism
Low	Low Or Normal	Low Or Normal	Nonthyroidal illness; pituitary (secondary) hypothyroidism
Normal	High	High	Thyroid hormone resistance syndrome (a mutation in the thyroid hormone receptor decreases thyroid hormone function)

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Urine Routine and Microscopic Examination

Physical Examination			
Volume *	20	ml	-
Colour *	Pale yellow	-	Pale yellow
Transparency *	Clear	-	Clear
Deposit *	Absent	-	Absent
Chemical Examination			
Reaction (pH) <i>Double Indicator</i>	6.0	-	4.5 - 8.0
Specific Gravity <i>Ion Exchange</i>	1.010	-	1.010 - 1.030
Urine Glucose (sugar) <i>Oxidase / Peroxidase</i>	Negative	-	Negative
Urine Protein (Albumin) <i>Acid / Base Colour Exchange</i>	Negative	-	Negative
Urine Ketones (Acetone) <i>Legals Test</i>	Negative	-	Negative
Blood <i>Peroxidase Hemoglobin</i>	Negative	-	Negative
Leucocyte esterase <i>Enzymatic Reaction</i>	Negative	-	Negative
Bilirubin Urine <i>Coupling Reaction</i>	Negative	-	Negative
Nitrite <i>Griless Test</i>	Negative	-	Negative
Urobilinogen <i>Ehrlichs Test</i>	Normal	-	Normal
Microscopic Examination			
Pus Cells (WBCs) *	1-2	/hpf	0 - 5
Epithelial Cells *	1-2	/hpf	0 - 4
Red blood Cells *	Absent	/hpf	Absent
Crystals *	Absent	-	Absent
Cast *	Absent	-	Absent
Yeast Cells *	Absent	-	Absent
Amorphous deposits *	Absent	-	Absent
Bacteria *	Absent	-	Absent
Protozoa *	Absent	-	Absent

Interpretation:

URINALYSIS- Routine urine analysis assists in screening and diagnosis of various metabolic, urological, kidney and liver disorders.

Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria, dehydration, urinary tract infections and acute illness with fever

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
<p>Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine. Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.</p>			
<p>Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine. Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise.</p>			
<p>Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders.</p>			
<p>Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys. Most common cause is bacterial urinary tract infection.</p>			
<p>Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration during infection increases with length of time the urine specimen is retained in bladder prior to collection.</p>			
<p>pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/ alkalosis or ingestion of certain type of food can affect the pH of urine.</p>			
<p>Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased specific gravity is seen in conditions like dehydration, glycosuria and proteinuria while decreased specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus.</p>			
<p>Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.</p>			
<p>Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in cases of haemolytic anaemia.</p>			

*** End Of Report ***

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.




Dr. Pragyamita Datta
MBBS, MD (Pathology)
Consultant Pathologist

ADPL HbA1c Graph Report

Name :	Sample Id :
Sample Type : Whole Blood EDTA	Total Area : 10.9

Peak Name	Retention Time(s)	Absorbance	Area	Result (Area %)
HbA0	67	0.1753	10.022	92.4
HbA1c	36	0.0051	0.587	5
La1c	34	0.0054	0.121	1.1
HbF	22	0.0013	0.068	0.6
Hba1b	16	0.002	0.078	0.7
Hba1a	12	0.0007	0.024	0.2



Terms and Conditions of Reporting

1. The presented findings in the Reports are intended solely for informational and interpretational purposes by the referring physician or other qualified medical professionals possessing a comprehensive understanding of reporting units, reference ranges, and technological limitations. The laboratory shall not be held liable for any interpretation or misinterpretation of the results, nor for any consequential or incidental damages arising from such interpretation.
2. It is to be presumed that the tests performed pertain to the specimen/sample attributed to the Customer's name or identification. It is presumed that the verification particulars have been cleared out by the customer or his/her representation at the point of generation of said specimen / sample. It is hereby clarified that the reports furnished are restricted solely to the given specimen only.
3. It is to be noted that variations in results may occur between different laboratories and over time, even for the same parameter for the same Customer. The assays are performed and conducted in accordance with standard procedures, and the reported outcomes are contingent on the specific individual assay methods and equipment(s) used, as well as the quality of the received specimen.
4. This report shall not be deemed valid or admissible for any medico-legal purposes.
5. The Customers assume full responsibility for apprising the Company of any factors that may impact the test finding. These factors, among others, includes dietary intake, alcohol, or medication / drug(s) consumption, or fasting. This list of factors is only representative and not exhaustive.

DISCLAIMER

This is a sample report provided for demonstration purposes only and does not represent an actual patient report. Test results, reference ranges, methodologies, instrumentation, and report formats may vary depending on the laboratory performing the test. The format and representation shown are indicative of reports generated by the National Reference Laboratory of Redcliffe Labs, Noida. This sample report should not be used for medical interpretation, diagnosis, or treatment decisions.