

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Extractable Nuclear Antigen (ENA) / ANA Panel, Qualitative (17 Antigens)

Mi-2 <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
Ku <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
U1-nRNP/Sm <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
Sm <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
SS-A(RO) <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
Ro- 52 <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
SS-B/La <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
Scl - 70 <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
PM-Scl <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
Jo - 1 <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
CENP-B <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
PCNA <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
ds-DNA <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
Nucleosomes <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
Histones <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
Rib- P Protein <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative
AMA- M2 <i>Immunoblot (Sample screening dilution is 1:101)</i>	Negative		Negative

Interpretation:

Test Description	Disease association
Mi-2	Dermatomyositis(15-30%) Idiopathic myositis(8-12 %)
Ku	SLE/SS(10-40 %) Myositis(40%)
U1-nRNP/Sm	Sharp syndrome(95%) SLE (15 - 40%) Poly-/dermatomyosities (12-16%) SS (2-12%)
Sm	SLE (15- 40%) - highly specific (> 90%)
SS-A (RO)	Sjogren syndrome(40-95%),SLE (20-60%)

Nitin

Dr. Nitin Arora
MBBS,MD (Microbiology)

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., H-55, Sector-63, Noida, Uttar Pradesh - 201301

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			

Test Description	Value(s)	Unit(s)	Reference Range
Ro 52	Neonatal lupus erythematosus NLE (95-100%)		
SS-B (La)	Sjogren syndrome (40-90%), SLE (10-20%) NLE(75%)		
Scl-70	SS(25 - 75%) - diffuse form (40 - 65%), & limited form (5 - 15%)		
PM-Sc	SS including overlap syndrome (10-20%) polymyositis(18%)		
Jo-1	Polymyositis/dermatomyositis (25-35%)		
CENP B, Centromeres	SS, limited from 80 - 95%,& diffuse form (8%) Primary biliary cirrhosis (10 - 30%)		
PCNA	SLE(3%)		
Double-stranded DNA	SLE (40-90%)		
Nucleosomes	SLE(40-70%)		
Histones	Drug induced LE (95 - 100%), SLE(50%) Rheumatoid arthritis (15 - 50%)		
Ribosomal PProtein	Drug induced LE (95 - 100%), SLE(50%) Rheumatoid arthritis (15 - 50%)		
AMA-M2 (Mitochondrial)	Primary biliary cirrhosis (>90%)		

COMMENTS


Immunoblot assay detects selected 17 ANAs which are most important & clinically relevant. However, in general, ANA includes many autoantibodies directed towards many nuclear(DNA & nucleoplasm) & cytoplasmic antigens, which are maximally screened & detected by using Hep-2 cells in indirect immunofluorescence method, but, not all of these are always clinically relevant antibodies. Hence, there is a possibility of patients having ANA positive by indirect immunofluorescence method but negative results on immunoblot. Also note that Immunoblot assay is more sensitive for Ro52/SSa , Scl 70,while poorly sensitive for DsDNA,hence such patients require further follow-up with nonspecific ELISAs based on clinical correlation for exact categorisation & diagnosis.

In case of weak Positive results repeat testing after 4-6 weeks or further testing with Monospecific Nuclear Antigens or Panels for confirmation of specific Autoantibodies is suggested.

Reference

Correlation of ANA by IIF patterns with immune profile using line immunoassay in indian scenario.Wendy Sebastian et al.Indian Juornal of Pathology and Microbiology-53(3),July-Sep 2010.

*** End Of Report ***



Dr. Nitin Arora
 MBBS,MD (Microbiology)

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., H-55, Sector-63, Noida, Uttar Pradesh - 201301

Terms and Conditions of Reporting

1. The presented findings in the Reports are intended solely for informational and interpretational purposes by the referring physician or other qualified medical professionals possessing a comprehensive understanding of reporting units, reference ranges, and technological limitations. The laboratory shall not be held liable for any interpretation or misinterpretation of the results, nor for any consequential or incidental damages arising from such interpretation.
2. It is to be presumed that the tests performed pertain to the specimen/sample attributed to the Customer's name or identification. It is presumed that the verification particulars have been cleared out by the customer or his/her representation at the point of generation of said specimen / sample. It is hereby clarified that the reports furnished are restricted solely to the given specimen only.
3. It is to be noted that variations in results may occur between different laboratories and over time, even for the same parameter for the same Customer. The assays are performed and conducted in accordance with standard procedures, and the reported outcomes are contingent on the specific individual assay methods and equipment(s) used, as well as the quality of the received specimen.
4. This report shall not be deemed valid or admissible for any medico-legal purposes.
5. The Customers assume full responsibility for apprising the Company of any factors that may impact the test finding. These factors, among others, includes dietary intake, alcohol, or medication / drug(s) consumption, or fasting. This list of factors is only representative and not exhaustive.

DISCLAIMER

This is a sample report provided for demonstration purposes only and does not represent an actual patient report. Test results, reference ranges, methodologies, instrumentation, and report formats may vary depending on the laboratory performing the test. The format and representation shown are indicative of reports generated by the National Reference Laboratory of Redcliffe Labs, Noida. This sample report should not be used for medical interpretation, diagnosis, or treatment decisions.