

smart Health Report

An Insightful Health Analytics Report
for Easier Understanding

Prepared For



Name

Gender

Patient ID

Age

Your Health at a Glance – A Personalized Journey

Report Sections

1

Body Summary

A visual snapshot of your overall health, simple and easy to understand

2

Quick Health Highlights

Your health scores and a single view of all abnormal results for quick attention

3

Lab Report Overview

Understand at a glance which tests are normal and which are abnormal

4

Personalized Health Advisory

Actionable insights and expert guidance tailored just for you

5

Doctor's Reference Report

Complete lab results with interpretations to share with your healthcare provider

How to Read This Report


This comprehensive health report provides detailed insights into your test results. Each section offers different perspectives on your health status, from visual summaries to detailed analysis and personalized recommendations.

Name Gender

Patient ID Age

● All In Range ● Borderline ● Out Of Range

Health Summary



Name Gender
Patient ID Age


Quick Health Summary

Personal Insights - Health Score

79

Overall, most parameters are within normal ranges, indicating good general health. The Thyroid may affect energy levels, and the Liver could influence overall vitality, so consider regular check-ups to monitor these areas. Adopting a balanced diet rich in fruits, vegetables, and juices, along with routine exercise like walking or yoga, can support your well-being. Remember, small lifestyle changes can lead to meaningful improvements in your health and vitality.

Note - Higher scores tentatively indicate better health status



Summary of Key Health Indicators

Total Parameters Tested	Borderline Results	Out Of Range Results
88	7	3

Health Status by Body System

Profile	Total	Borderline	Out of Range	Key Results
Inflammation	1	0	1	● ESR (23)
Blood Disorder	17	3	1	● Abs. Basophil Count (0) ● Haemoglobin (11.7) ● MPV (14.4)
Liver Profile	15	0	1	● GGT (4.7)
Vitamin Profile	1	0	0	All In Range
Thyroid Profile	3	1	0	● TSH (5.78)
Anemia Studies	8	0	0	All In Range
Infectious Diseases	6	0	0	All In Range
Diabetes Monitoring	2	0	0	All In Range
Kidney Profile	10	2	0	● BUN : Creatinine ratio (20.63) ● Urea : Creatinine ratio (44.17)
Urinalysis	12	0	0	All In Range
Electrolytes	4	0	0	All In Range
Cardiac Profile	9	1	0	● Total Cholesterol : HDL ratio (3.08)

Name Gender

Patient ID Age

Report Summary ● In Range ● Borderline ● Out Of Range ● No color - Reference range not available

INFLAMMATION		
Test Name	Result unit	Range
● ESR - Erythrocyte Sedimentation Rate	23 mm/hr	< 12

VITAMIN PROFILE		
Test Name	Result unit	Range
● Vitamin D 25 - Hydroxy	33.1 ng/mL	30 - 100

THYROID PROFILE		
Test Name	Result unit	Range
● Triiodothyronine (T3)	139 ng/dL	35 - 193
● Total Thyroxine (T4)	8.38 µg/dL	4.87 - 11.2
● Thyroid Stimulating Hormone (Ultrasensitive)	5.78 µIU/mL	0.35 - 4.94

BLOOD DISORDER		
Test Name	Result unit	Range
● Hemoglobin	11.7 g/dL	12 - 15
● TLC	6.6 $10^3/\mu\text{l}$	4 - 10
● Neutrophils	62 %	40 - 80
● Lymphocytes	32 %	20 - 40
● Monocytes	4 %	2 - 10
● Eosinophils	2 %	1 - 6
● Basophils	0 %	< 2
● Neutrophils.	4.09 $10^3/\mu\text{l}$	2 - 7
● Lymphocytes.	2.11 $10^3/\mu\text{l}$	1 - 3
● Monocytes.	0.26 $10^3/\mu\text{l}$	0.2 - 1
● Eosinophils.	0.13 $10^3/\mu\text{l}$	0.02 - 0.5
● Basophils.	0 $10^3/\mu\text{l}$	0.02 - 0.5
● Platelet Count	160 $10^3/\mu\text{l}$	150 - 410
● Mean Platelet Volume (MPV)	14.4 fL	9.3 - 12.1
● PDW	22 fL	8.3 - 25
● P-LCR	55.5 %	18 - 50
● P-LCC	78 $10^9/L$	44 - 140

Name Gender

Patient ID Age

Report Summary ● In Range ● Borderline ● Out Of Range ● No color - Reference range not available

ANEMIA STUDIES

Test Name	Result <small>unit</small>	Range
● RBC Count	4.1 $10^6/\mu\text{L}$	3.8 - 4.8
● PCV	36 %	36 - 46
● MCV	87 fl	83 - 101
● MCH	28.2 pg	27 - 32
● MCHC	32.4 g/dL	31.5 - 34.5
● RDW (CV)	13.6 %	11.6 - 14
● RDW-SD	42.1 fl	35.1 - 43.9
Mentzer Index	21.22 %	

INFECTIOUS DISEASES

Test Name	Result <small>unit</small>	Range
● PCT	0.2 %	0.17 - 0.32
Deposit	Absent	
Leucocyte esterase	Negative	
Pus Cells (WBCs)	3-4 /hpf	
Yeast Cells	Absent	
Protozoa	Absent	

DIABETES MONITORING

Test Name	Result <small>unit</small>	Range
● Glucose Fasting	94.3 mg/dL	70 - 100
Urine Glucose (sugar)	Negative	

Name Gender

Patient ID Age

Report Summary ● In Range ● Borderline ● Out Of Range ● No color - Reference range not available

LIVER PROFILE		
Test Name	Result <small>unit</small>	Range
● Bilirubin Total	0.4 mg/dL	0.2 - 1.2
● Bilirubin Direct	0.2 mg/dL	< 0.5
● Bilirubin Indirect	0.2 mg/dL	0.1 - 1
● SGOT/AST	23.9 U/L	5 - 34
● SGPT/ALT	22.7 U/L	< 55
SGOT/SGPT Ratio	1.05 %	
● Alkaline Phosphatase	131 U/L	40 - 150
● Total Protein	7 g/dL	6.4 - 8.3
● Albumin	4.2 gm/dL	3.8 - 5
● Globulin	2.8 g/dL	2.3 - 3.5
● Albumin :Globulin Ratio	1.5	1 - 2.1
● Gamma Glutamyl Transferase (GGT)	4.7 U/L	9 - 36
● Calcium Serum	9.9 mg/dL	8.4 - 10.2
Bilirubin Urine	Negative	
Urobilinogen	Normal	

KIDNEY PROFILE		
Test Name	Result <small>unit</small>	Range
● Blood Urea	26.5 mg/dL	19 - 44.1
● Bun	12.38 mg/dL	6 - 20
● Creatinine	0.6 mg/dL	0.57 - 1.11
eGFR (CKD-EPI)	114.12 ml/min/1.73 sq m	
● Bun/Creatinine Ratio	20.63	12 - 20
● Urea / Creatinine Ratio	44.17	25.68 - 42.8
Urine Protein (Albumin)	Negative	
Blood	Negative	
Crystals	Absent	
Cast	Absent	

Name Gender

Patient ID Age

Report Summary

● In Range ● Borderline ● Out Of Range No color - Reference range not available

URINALYSIS

Test Name	Result unit	Range
● Uric Acid	4.6 mg/dL	2.6 - 6
Volume	15 mL	
Colour	Pale yellow	
Transparency	Clear	
● Reaction (pH)	5.0	4.5 - 8
● Specific Gravity	1.010	1.01 - 1.03
Urine Ketones (Acetone)	Negative	
Nitrite	Negative	
Epithelial Cells	1-3 /hpf	
Red blood Cells	Absent /hpf	
Amorphous deposits	Absent	
Bacteria	Absent	

ELECTROLYTE PROFILE

Test Name	Result unit	Range
● Phosphorus	3.5 mg/dL	2.3 - 4.7
● Sodium	139.7 mmol/L	136 - 145
● Potassium	5.0 mmol/L	3.5 - 5.1
● Chloride	105.6 mmol/L	98 - 107

CARDIAC PROFILE

Test Name	Result unit	Range
● Total Cholesterol	151 mg/dL	< 200
● Triglycerides	74.9 mg/dL	< 150
● HDL Cholesterol	49 mg/dL	40 - 80
● Non HDL Cholesterol	102 mg/dL	< 130
● LDL Cholesterol	87.02 mg/dL	30 - 100
● V.L.D.L Cholesterol	14.98 mg/dL	< 30
● Chol/HDL Ratio	3.08 Ratio	3.5 - 5
● HDL/ LDL Ratio	0.56 Ratio	0.5 - 3
LDL/HDL Ratio	1.78 Ratio	

Name

Gender

Patient ID

Age

Health Advisory

● In Range ● Borderline (BL) ● Out Of Range



Inflammation

Inflammation is the body's immune system's response to an injury, surgery, or irritation. This natural defense process acts by removing injurious stimuli and initiating the healing process. Inflammation can be chronic (such as arthritis) or acute (like in case of trauma).

ESR - Erythrocyte Sedimentation Rate: 23 mm/hr

● OUT OF RANGE



Thyroid

This panel is used to check the imbalance in your thyroid gland. A healthy thyroid gland is very important for metabolism, controlling body temperature, regulation of mood, muscle strength and regulation of body weight

Thyroid Stimulating Hormone (Ultrasensitive): 5.78 μ U/mL

● BORDERLINE



Blood Disorder

Blood disorders affect one or more components of blood such as red blood cells, white blood cells, platelets, or plasma. These tests help in diagnosing conditions like anemia, clotting disorders, infections, and other hematological abnormalities.

Hemoglobin: 11.7 g/dL

● BORDERLINE





Liver Profile

One of the main functions of your liver is to make proteins that are secreted in your blood. It also makes enzymes which convert food into energy, and processes old muscles and cells. When your liver is damaged, enzymes leak into your blood and appear in the blood test

Gamma Glutamyl Transferase (GGT): 4.7 U/L

● OUT OF RANGE



Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			
Test Description	Value(s)	Unit(s)	Reference Range

Smart Full Body Check up with Vitamin D

Complete Blood Count (CBC)

RBC Parameters			
Hemoglobin <i>colorimetric</i>	11.7 L*	g/dL	12.0-15.0
RBC Count <i>Electrical impedance</i>	4.1	10 ⁶ /μl	3.8 - 4.8
PCV <i>Calculated</i>	36	%	36 - 46
MCV <i>Calculated</i>	87	fl	83 - 101
MCH <i>Calculated</i>	28.2	pg	27 - 32
MCHC <i>Calculated</i>	32.4	g/dL	31.5 - 34.5
RDW (CV) * <i>Calculated</i>	13.6	%	11.6 - 14.0
RDW-SD * <i>Calculated</i>	42.1	fl	35.1 - 43.9
WBC Parameters			
TLC <i>Electrical impedance and microscopy</i>	6.6	10 ³ /μl	4 - 10
Differential Leucocyte Count			
Neutrophils	62	%	40-80
Lymphocytes	32	%	20-40
Monocytes	4	%	2-10
Eosinophils	2	%	1-6
Basophils	0	%	<2
Absolute Leukocyte Counts			
Neutrophils.	4.09	10 ³ /μl	2 - 7
Lymphocytes.	2.11	10 ³ /μl	1 - 3
Monocytes.	0.26	10 ³ /μl	0.2 - 1.0
Eosinophils.	0.13	10 ³ /μl	0.02 - 0.5
Basophils.	0	10 ³ /μl	0.02 - 0.5
Platelet Parameters			
Platelet Count <i>Electrical impedance and microscopy</i>	160	10 ³ /μl	150 - 410
Mean Platelet Volume (MPV) * <i>Calculated</i>	14.4 H*	fL	9.3 - 12.1
PCT * <i>Calculated</i>	0.2	%	0.17 - 0.32

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



DR. RAGINI GUPTA
MD PATHOLOGY
CONSULTANT PATHOLOGIST

Patient NAME :		Report STATUS
DOB/Age/Gender :		Barcode NO
Patient ID / UHID :		Sample Type
Referred BY :		Report Date
Sample Collected :		

Test Description	Value(s)	Unit(s)	Reference Range
PDW * <i>Calculated</i>	22	fL	8.3 - 25.0
P-LCR * <i>Calculated</i>	55.5 H*	%	18 - 50
P-LCC * <i>Calculated</i>	78	10 ⁹ /L	44 - 140
Mentzer Index * <i>Calculated</i>	21.22	%	> 13

Interpretation:

CBC provides information about red cells, white cells and platelets. Results are useful in the diagnosis of anemia, infections, leukemias, clotting disorders and many other medical conditions.

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



DR. RAGINI GUPTA
MD PATHOLOGY
CONSULTANT PATHOLOGIST

Patient NAME :		Report STATUS :	
DOB/Age/Gender :		Barcode NO :	
Patient ID / UHID :		Sample Type :	
Referred BY :		Report Date :	
Sample Collected :			
Test Description	Value(s)	Unit(s)	Reference Range

Erythrocyte Sedimentation Rate (ESR)

ESR - Erythrocyte Sedimentation Rate <i>MODIFIED WESTERGREN</i>	23 H*	mm/hr	0 - 12
--	--------------	-------	--------

Interpretation:

ESR is also known as Erythrocyte Sedimentation Rate. An ESR test is used to assess inflammation in the body. Many conditions can cause an abnormal ESR, so an ESR test is typically used with other tests to diagnose and monitor different diseases. An elevated ESR may occur in inflammatory conditions including infection, rheumatoid arthritis, systemic vasculitis, anemia, multiple myeloma, etc. Low levels are typically seen in congestive heart failure, polycythemia, sickle cell anemia, hypo fibrinogenemia, etc.

Reference- Dacie and Lewis practical hematology

Note :- (H* - High, L* - Low, CL* - Critical Low, CH* - Critical High)

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



DR. RAGINI GUPTA
MD PATHOLOGY
CONSULTANT PATHOLOGIST

Patient NAME		Report STATUS :	
DOB/Age/Gender		Barcode NO :	
Patient ID / UHID		Sample Type :	
Referred BY		Report Date :	
Sample Collected			
Test Description	Value(s)	Unit(s)	Reference Range

Glucose Fasting

Glucose Fasting <i>Hexokinase</i>	94.3	mg/dL	70 - 100
--------------------------------------	------	-------	----------

Interpretation:

Status	Fasting plasma glucose in mg/dL
Normal	70 - 100
Impaired fasting glucose	101 - 125
Diabetes	≥126

Reference : American Diabetes Association

Comment :

Blood glucose determinations are commonly used as an aid in the diagnosis and treatment of diabetes. Elevated glucose levels (hyperglycemia) may also occur with pancreatic neoplasm, hyperthyroidism, and adrenal cortical hyper function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy insulinoma, or various liver diseases.

Note

- 1.The diagnosis of Diabetes requires a fasting plasma glucose of > or = 126 mg/dL or a random / 2 hour plasma glucose value of > or = 200 mg/dL with symptoms of diabetes mellitus.
- 2.Very high glucose levels (>450 mg/dL in adults) may result in Diabetic Ketoacidosis.

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



DR. RAGINI GUPTA
MD PATHOLOGY
CONSULTANT PATHOLOGIST

Patient NAME :		Report STATUS	
DOB/Age/Gender :		Barcode NO	
Patient ID / UHID :		Sample Type	
Referred BY :		Report Date	
Sample Collected :			
Test Description	Value(s)	Unit(s)	Reference Range

Liver Function Test (LFT)

Bilirubin Total <i>Photometric</i>	0.4	mg/dL	0.2 - 1.2
Bilirubin Direct * <i>Diazo Reaction</i>	0.2	mg/dL	0.0 - 0.5
Bilirubin Indirect * <i>Calculation (T Bil - D Bil)</i>	0.2	mg/dL	0.1 - 1.0
SGOT/AST <i>IFCC without P5P</i>	23.9	U/L	5 - 34
SGPT/ALT <i>IFCC without P5P</i>	22.7	U/L	0 to 55
SGOT/SGPT Ratio *	1.05	-	-
Alkaline Phosphatase <i>IFCC</i>	131	U/L	40 - 150
Total Protein <i>Biuret</i>	7	g/dL	6.4 - 8.3
Albumin <i>BCG</i>	4.2	gm/dL	3.8 - 5.0
Globulin * <i>Calculation (T.P - Albumin)</i>	2.8	g/dL	2.3 - 3.5
Albumin :Globulin Ratio * <i>Calculation (Albumin/Globulin)</i>	1.5	-	1.0 - 2.1
Gamma Glutamyl Transferase (GGT) * <i>Photometric</i>	4.7 L*	U/L	9 - 36

Interpretation:

The liver filters blood, metabolizes nutrients, detoxifies harmful substances, and produces blood clotting proteins. Liver cells contain enzymes that facilitate these functions. When cells are damaged, enzymes leak into the blood, detectable through blood tests.

Key enzymes tested:

- AST (SGOT):** may indicate tissue injury / damage in muscles or liver.
- ALT (SGPT):** Primarily in the liver. Elevated ALT and AST suggest liver damage.
- Alkaline Phosphatase & GGT:** Linked to bile production and flow. Elevated levels may indicate bile flow issues related to the liver, gallbladder, or bile ducts.

Blood proteins, **albumin and globulin**, are essential for growth, development, and health.

- Low protein:** May indicate bleeding, liver disorders, malnutrition, or agammaglobulinemia.
- High protein (Hyperproteinemia):** Often due to dehydration or increased protein production.
- Low albumin:** Caused by poor diet, kidney, or liver disease.
- High albumin:** Usually due to severe dehydration.

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



DR. RAGINI GUPTA
MD PATHOLOGY
CONSULTANT PATHOLOGIST

Patient NAME	
DOB/Age/Gender	Report STATUS
Patient ID / UHID	Barcode NO
Referred BY	Sample Type
Sample Collected	Report Date

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Kidney Function Test (KFT)

Blood Urea <i>Urease</i>	26.5	mg/dL	19 - 44.1
Bun * <i>Urease</i>	12.38	mg/dL	6 - 20
Creatinine <i>Photometric</i>	0.6	mg/dL	0.57 - 1.11
eGFR (CKD-EPI)	114.12	ml/min/1.73 sq m	Normal Or High: ≥ 90 Mild Or Decrease: 60-89 Mild To Moderate Decrease: 45-59 Mild To Severe Decrease: 30-44 Severe Decrease: 15-29 Kidney Failure: < 15
Bun/Creatinine Ratio * <i>Calculated</i>	20.63 H*		12 - 20
Urea / Creatinine Ratio * <i>Calculated</i>	44.17 H*		25.68- 42.8
Uric Acid <i>Uricase</i>	4.6	mg/dL	2.6 - 6.0
Calcium Serum <i>Arsenazo III</i>	9.9	mg/dL	8.4 - 10.2
Phosphorus <i>Photometric</i>	3.5	mg/dL	2.3 - 4.7
Sodium <i>Potentiometric</i>	139.7	mmol/L	136 - 145
Potassium <i>Potentiometric</i>	5.0	mmol/L	3.5 - 5.1
Chloride <i>Potentiometric</i>	105.6	mmol/L	98 - 107

Interpretation:

Kidney function tests is a collective term for a variety of individual tests and procedures that can be done to evaluate how well the kidneys are functioning. Many conditions can affect the ability of the kidneys to carry out their vital functions. Some lead to a rapid (acute) decline in kidney function others lead to a gradual (chronic) decline in function. Both result in a buildup of toxic waste substances done on urine samples, as well as on blood samples. A number of symptoms may indicate a problem with your kidneys. These include : high blood pressure, blood in urine, frequent urges to urinate, difficulty beginning urination, painful urination, swelling in the hands and feet due to a buildup of fluids in the body. A single symptom may not mean something serious. However, when occurring simultaneously, these symptoms suggest that your kidneys are not working properly. Kidney function tests can help determine the reason. Ionized calcium for this test if you have signs of kidney or parathyroid disease. The test may also be done to monitor progress and treatment of these diseases."eGFR test is applicable for patients aged 18 years or more."

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)

(*) Parameter(s) are outside the scope of tests recognized under the NABL M(EL)T Scheme.



DR. RAGINI GUPTA
MD PATHOLOGY
CONSULTANT PATHOLOGIST

Booking Centre :- Aradhya Collection Center, Basawan Road, Lalapur, Kudra, Kaimur, Bihar- 821101

Processing Lab :- Redcliffe Lifetech Pvt. Ltd., House No. 59, south opposite of Block-1, madhuban Apartment Kailashpuri, Malahi Pakri, kankarbagh, Behind raymond showroom, Patna, Bihar - 800020. "

📞 928-909-0609

✉️ ccsupport@redcliffelabs.com

🌐 www.redcliffelabs.com

All Lab results are subject to clinical interpretation by qualified medical professional and this report is not subject to use for any medico-legal purpose.

Patient NAME		Report STATUS :	
DOB/Age/Gender		Barcode NO :	
Patient ID / UHID		Sample Type :	
Referred BY		Report Date :	
Sample Collected			

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Lipid Profile

Total Cholesterol <i>Enzymatic - Cholesterol Oxidase</i>	151	mg/dL	<200
Triglycerides <i>Colorimetric - Lip/Glycerol Kinase</i>	74.9	mg/dL	<150
HDL Cholesterol <i>Accelerator Selective Detergent</i>	49	mg/dL	>40
Non HDL Cholesterol * <i>Calculated</i>	102	mg/dL	<130
LDL Cholesterol * <i>Calculated</i>	87.02	mg/dL	<100
V.L.D.L Cholesterol * <i>Calculated</i>	14.98	mg/dL	< 30
Chol/HDL Ratio * <i>Calculated</i>	3.08 L*	Ratio	3.5 - 5.0
HDL/ LDL Ratio * <i>Calculated</i>	0.56	Ratio	0.5 - 3.0
LDL/HDL Ratio * <i>Calculated</i>	1.78	Ratio	-

Interpretation:

Lipid level assessments must be made following 9 to 12 hours of fasting, otherwise assay results might lead to erroneous interpretation. NCEP recommends of 3 different samples to be drawn at intervals of 1 week for harmonizing biological variables that might be encountered in single assays.

National Lipid Association Recommendations (NLA-2014)	Total Cholesterol (mg/dL)	Triglyceride (mg/dL)	LDL Cholesterol (mg/dL)	Non HDL Cholesterol (mg/dL)
Optimal	<200	<150	<100	<130
Above Optimal			100-129	130 - 159
Borderline High	200-239	150-199	130-159	160 - 189
High	>=240	200-499	160-189	190 - 219
Very High	-	>=500	>=190	>=220

HDL Cholesterol	
Low	High
<40	>=60

Risk Stratification for ASCVD (Atherosclerotic Cardiovascular Disease) by Lipid Association of India.

Risk Category	A. CAD with > 1 feature of high risk group
Extreme risk group	B. CAD with >1 feature of very high risk group of recurrent ACS (within 1 year) despite LDL-C <or = 50 mg/dl or poly vascular disease
Very High Risk	1.Established ASCVD 2.Diabetes with 2 major risk factors of evidence of end organ damage 3. Familial Homozygous Hypercholesterolemia
	1. Three major ASCVD risk factors 2. Diabetes with 1 major risk factor or no evidence

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)



DR. RAGINI GUPTA
MD PATHOLOGY
CONSULTANT PATHOLOGIST

Patient NAME :
 DOB/Age/Gender : Report STATUS
 Patient ID / UHID : Barcode NO
 Referred BY : Sample Type
 Sample Collected : Report Date

Test Description	Value(s)	Unit(s)	Reference Range
High Risk	of end organ damage 3. CHD stage 3B or 4. 4 LDL >190 mg/dl 5. Extreme of a single risk factor 6. Coronary Artery Calcium - CAC > 300 AU 7. Lipoprotein a >= 50 mg/dl 8. Non stenotic carotid plaque		
Moderate Risk	2 major ASCVD risk factors		
Low Risk	0-1 major ASCVD risk factors		
Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors			
1. Age >=45 years in Males & >= 55 years in Females	3. Current Cigarette smoking or tobacco use		
2. Family history of premature ASCVD	4. High blood pressure		
5. Low HDL			

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by Lipid Association of India in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group Category A	<50 (Optional goal <OR = 30)	<80 (Optional goal <OR = 60)	>OR = 50	>OR = 80
Extreme Risk Group Category B	>OR = 30	>OR = 60	> 30	> 60
Very High Risk	<50	<80	>OR = 50	>OR = 80
High Risk	<70	<100	>OR = 70	>OR = 100
Moderate Risk	<100	<130	>OR = 100	>OR = 130
Low Risk	<100	<130	>OR = 130*	>OR = 160

* After an adequate non-pharmacological intervention for at least 3 months.

References : Management of Dyslipidaemia for the Prevention of Stroke : Clinical practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology,2022,20,134-155.

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)



DR. RAGINI GUPTA
MD PATHOLOGY
CONSULTANT PATHOLOGIST

Patient NAME		Report STATUS	
DOB/Age/Gender		Barcode NO	
Patient ID / UHID		Sample Type	
Referred BY		Report Date	
Sample Collected			
Test Description	Value(s)	Unit(s)	Reference Range

Vitamin D 25 Hydroxy

Vitamin D 25 - Hydroxy <i>CMIA</i>	33.1	ng/mL	Deficiency : <30 ng/mL
---------------------------------------	------	-------	------------------------

Interpretation:

25-Hydroxy vitamin D represents the main body reservoir and transport form. Mild to moderate deficiency is associated with Osteoporosis / Secondary Hyperparathyroidism while severe deficiency causes Rickets in children and Osteomalacia in adults. Prevalence of Vitamin D deficiency is approximately >50% specially in the elderly. This assay is useful for diagnosis of vitamin D deficiency and Hypervitaminosis D. It is also used for differential diagnosis of causes of Rickets & Osteomalacia and for monitoring Vitamin D replacement therapy.



DR. RAGINI GUPTA
MD PATHOLOGY
CONSULTANT PATHOLOGIST

Patient NAME :
 DOB/Age/Gender :
 Patient ID / UHID :
 Referred BY :
 Sample Collected :

Report STATUS :
 Barcode NO :
 Sample Type :
 Report Date :

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Thyroid Profile Total

Triiodothyronine (T3) CMIA	139	ng/dL	35 - 193
Total Thyroxine (T4) CMIA	8.38	µg/dL	4.87 - 11.2
Thyroid Stimulating Hormone (Ultrasensitive) CMIA	5.78 H*	µIU/mL	0.35 - 4.94

Interpretation:

Pregnancy	Reference Range TSH
1st Trimester	0.1 - 2.5
2nd Trimester	0.2 - 3.0
3rd Trimester	0.3 - 3.0

Clinical Use:

1. Diagnose Hypothyroidism & Hyperthyroidism
2. Monitor T4 therapy
3. Measure subnormal TSH levels

Increased TSH: Primary hypothyroidism, Subclinical hypothyroidism, TSH-dependent hyperthyroidism, Thyroid hormone resistance

Decreased TSH: Graves' disease, Autonomous thyroid hormone secretion, TSH deficiency

Thyroid malfunction (hyper or hypo) affects T3 & T4 levels. Pituitary or hypothalamic issues also influence thyroid activity.

1. **Primary Hypothyroidism:** High TSH levels.
2. **Secondary/Tertiary Hypothyroidism:** Low TSH levels.
3. **Euthyroid Sick Syndrome:** Abnormal thyroid test results due to non-thyroidal illnesses (NTI).

TBG levels are stable in healthy individuals but may be altered by pregnancy, estrogens, androgens, steroids, or glucocorticoids, causing inaccurate T3 & T4 readings.

TSH	T4	T3	Interpretation
High	Normal	Normal	Mild (subclinical) hypothyroidism
High	Low	Low Or Normal	Hypothyroidism
Low	Normal	Normal	Mild (subclinical) hyperthyroidism
Low	High Or Normal	High Or Normal	Hyperthyroidism
Low	Low Or Normal	Low Or Normal	Nonthyroidal illness; pituitary (secondary) hypothyroidism
Normal	High	High	Thyroid hormone resistance syndrome (a mutation in the thyroid hormone receptor decreases thyroid hormone function)

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)



DR. RAGINI GUPTA
 MD PATHOLOGY
 CONSULTANT PATHOLOGIST

Patient NAME	Report STATUS :
DOB/Age/Gender	Barcode NO :
Patient ID / UHID	Sample Type :
Referred BY	Report Date :
Sample Collected	

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Note :- (H* - High , L* - Low ,CL* - Critical Low,CH* - Critical High)



DR. RAGINI GUPTA
MD PATHOLOGY
CONSULTANT PATHOLOGIST

Patient NAME	Report STATUS :
DOB/Age/Gender	Barcode NO :
Patient ID / UHID	Sample Type :
Referred BY	Report Date :
Sample Collected	

Test Description	Value(s)	Unit(s)	Reference Range
------------------	----------	---------	-----------------

Urine Routine and Microscopic Examination

Physical Examination			
Volume *	15	mL	-
Colour *	Pale yellow	-	Pale yellow
Transparency *	Clear	-	Clear
Deposit *	Absent	-	Absent
Chemical Examination			
Reaction (pH) <i>Double Indicator</i>	5.0	-	4.5 - 8.0
Specific Gravity <i>Ion Exchange</i>	1.010	-	1.010 - 1.030
Urine Glucose (sugar) <i>Oxidase / Peroxidase</i>	Negative	-	Negative
Urine Protein (Albumin) <i>Acid / Base Colour Exchange</i>	Negative	-	Negative
Urine Ketones (Acetone) <i>Legals Test</i>	Negative	-	Negative
Blood <i>Peroxidase Hemoglobin</i>	Negative	-	Negative
Leucocyte esterase <i>Enzymatic Reaction</i>	Negative	-	Negative
Bilirubin Urine <i>Coupling Reaction</i>	Negative	-	Negative
Nitrite <i>Griless Test</i>	Negative	-	Negative
Urobilinogen <i>Ehrlichs Test</i>	Normal	-	Normal
Microscopic Examination			
Pus Cells (WBCs) *	3-4	/hpf	0 - 5
Epithelial Cells *	1-3	/hpf	0 - 4
Red blood Cells *	Absent	/hpf	Absent
Crystals *	Absent	-	Absent
Cast *	Absent	-	Absent
Yeast Cells *	Absent	-	Absent
Amorphous deposits *	Absent	-	Absent
Bacteria	Absent	-	Absent
Protozoa *	Absent	-	Absent

Interpretation:

URINALYSIS- Routine urine analysis assists in screening and diagnosis of various metabolic, urological, kidney and liver disorders.

Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria, dehydration, urinary tract infections and acute illness with fever



DR. RAGINI GUPTA
MD PATHOLOGY
CONSULTANT PATHOLOGIST

Patient NAME		Report STATUS :
DOB/Age/Gender		Barcode NO :
Patient ID / UHID		Sample Type :
Referred BY		Report Date :
Sample Collected		

Test Description	Value(s)	Unit(s)	Reference Range
<p>Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine. Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.</p>			
<p>Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine. Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise.</p>			
<p>Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders.</p>			
<p>Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys. Most common cause is bacterial urinary tract infection.</p>			
<p>Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration during infection increases with length of time the urine specimen is retained in bladder prior to collection.</p>			
<p>pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/ alkalosis or ingestion of certain type of food can affect the pH of urine.</p>			
<p>Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased specific gravity is seen in conditions like dehydration, glycosuria and proteinuria while decreased specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus.</p>			
<p>Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.</p>			
<p>Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in cases of haemolytic anaemia.</p>			

*** End Of Report ***



DR. RAGINI GUPTA
MD PATHOLOGY
CONSULTANT PATHOLOGIST

Terms and Conditions of Reporting

1. The presented findings in the Reports are intended solely for informational and interpretational purposes by the referring physician or other qualified medical professionals possessing a comprehensive understanding of reporting units, reference ranges, and technological limitations. The laboratory shall not be held liable for any interpretation or misinterpretation of the results, nor for any consequential or incidental damages arising from such interpretation.
2. It is to be presumed that the tests performed pertain to the specimen/sample attributed to the Customer's name or identification. It is presumed that the verification particulars have been cleared out by the customer or his/her representation at the point of generation of said specimen / sample. It is hereby clarified that the reports furnished are restricted solely to the given specimen only.
3. It is to be noted that variations in results may occur between different laboratories and over time, even for the same parameter for the same Customer. The assays are performed and conducted in accordance with standard procedures, and the reported outcomes are contingent on the specific individual assay methods and equipment(s) used, as well as the quality of the received specimen.
4. This report shall not be deemed valid or admissible for any medico-legal purposes.
5. The Customers assume full responsibility for apprising the Company of any factors that may impact the test finding. These factors, among others, includes dietary intake, alcohol, or medication / drug(s) consumption, or fasting. This list of factors is only representative and not exhaustive.

About Redcliffe Labs

We are India's Most Trusted & Fastest Growing Network of Diagnostics Labs

Best Customer Experience



Commitment to excellence, high end technology oriented staff

100% Report Correctness



Focus on quality with accurate results

Best Prices With Fast Reports



Value for money with quick turn around time (TAT)

Your booking just gave back to nature – with every health checkup, you're contributing by planting a tree!



BharatFit -5

BEST SELLER
★★★

₹2399 ~~₹4214~~

96 TEST PARAMETERS

- ✓ Blood Sugar Fasting (1 Test)
- ✓ Lipid Profile (9 Tests)
- ✓ Liver Function Test (12 Tests)
- ✓ Kidney Function Test (12 Tests)
- ✓ Thyroid Profile Total (3 Tests)
- ✓ Urine R/M (23 Tests)
- ✓ Complete Blood Count (26 Tests)
- ✓ ESR (1 Test)
- ✓ HbA1c (2 Tests)
- ✓ Vitamin D (1 Test)
- ✓ Vitamin B12 (1 Test)
- ✓ Iron Studies (4 Tests)
- ✓ HBsAg (Rapid) (1 Test)



3600 + Tests & Packages



220+ Cities Presence



Home collection available



100% Accurate Report Guarantee.



Terms and Conditions of Reporting

1. The presented findings in the Reports are intended solely for informational and interpretational purposes by the referring physician or other qualified medical professionals possessing a comprehensive understanding of reporting units, reference ranges, and technological limitations. The laboratory shall not be held liable for any interpretation or misinterpretation of the results, nor for any consequential or incidental damages arising from such interpretation.
2. It is to be presumed that the tests performed pertain to the specimen/sample attributed to the Customer's name or identification. It is presumed that the verification particulars have been cleared out by the customer or his/her representation at the point of generation of said specimen / sample. It is hereby clarified that the reports furnished are restricted solely to the given specimen only.
3. It is to be noted that variations in results may occur between different laboratories and over time, even for the same parameter for the same Customer. The assays are performed and conducted in accordance with standard procedures, and the reported outcomes are contingent on the specific individual assay methods and equipment(s) used, as well as the quality of the received specimen.
4. This report shall not be deemed valid or admissible for any medico-legal purposes.
5. The Customers assume full responsibility for apprising the Company of any factors that may impact the test finding. These factors, among others, includes dietary intake, alcohol, or medication / drug(s) consumption, or fasting. This list of factors is only representative and not exhaustive.

DISCLAIMER

This is a sample report provided for demonstration purposes only and does not represent an actual patient report. Test results, reference ranges, methodologies, instrumentation, and report formats may vary depending on the laboratory performing the test. The format and representation shown are indicative of reports generated by the National Reference Laboratory of Redcliffe Labs, Noida. This sample report should not be used for medical interpretation, diagnosis, or treatment decisions.